

Chemistry in Everyday Life


Drugs and Their Classification

Drugs

- Chemicals of low molecular masses ($\sim 100\text{--}500$ u) used for producing biological response by interacting with target macromolecules
- Drugs used for therapeutic effect are called medicines.
- Chemotherapy refers to the use of chemicals for therapeutic effect.

Classification of Drugs

- Based on pharmacological effect:
 - Useful for doctors as it provides the whole range of drugs available to cure a particular type of problem
- Based on drug action:
 - On the basis of the action of a drug on a particular biochemical process
- Based on chemical structure:
 - Drugs classified on this basis have common structural features, and also may have similar pharmacological activity. For example, sulphonamides have common structural features as follows:




- Based on molecular targets:
 - Useful for medicinal chemists
 - Target molecules or molecular targets are the biomolecules with which drugs interact; for example, carbohydrates, lipids, proteins and nucleic acids.


Drug–Target Interaction

Enzymes as drug targets


- Catalytic action of enzymes: Two major functions of enzymes are –
- Substrate molecules are held in suitable positions by the active sites of enzymes through ionic bonding, H-bonding, van der Waals interaction or dipole–dipole interaction.



- To attack the substrate molecules, functional groups are provided by enzymes.
- Drug–enzyme interaction
- Drugs inhibit the catalytic action of enzymes by blocking the binding site of enzymes. Such drugs are called enzyme inhibitors.
- The two ways of drug–enzyme interaction are:
- Drugs compete with substrate molecules to attack the active site of enzymes. Such drugs are called competitive inhibitors.



- Some drugs change the shape of the active site of an enzyme. Such drugs bind to a different site, which is called allosteric site, and changes the shape of the active site in such a way that the substrate cannot recognise it.



Receptors as drug targets

- Receptors are proteins that are crucial to the communication process of the body.
- Chemical messengers are the chemicals through which messages between two neurons and between neurons and muscles are communicated.
- Antagonists are drugs which bind to the receptor site and inhibit its natural function.
- Agonists are drugs that mimic the natural messenger by switching on the receptor.

Therapeutic Action of Different Classes of Drugs


Antacids

- Used for the treatment of acidity
- Example: Sodium hydrogencarbonate or a mixture of aluminium and magnesium hydroxide
- Metal hydroxides are better alternatives.

Reason: They do not increase the pH above neutrality as they are insoluble.

Anti-histamines


- **Histamines**
- Stimulate secretion of pepsin and hydrochloric acid in the stomach
- Also responsible for the nasal congestion associated with common cold, and allergic response to pollen




Histamine

- **Anti-histamine Drugs**

- Cimetidine and ranitidine prevent the interaction of histamine with the receptors present in the stomach walls, and as a result, lesser amount of acid is released.



Cimetidine




Ranitidine

- Brompheniramine (Dimetapp) and terfenadine (Seldane) interfere with the natural action of histamine by competing with histamine for binding sites of receptors.
- These anti-histamines do not affect the secretion of acid in the stomach. Reason: Anti-allergic and antacid drugs work on different receptors.



Brompheniramine
(Dimetapp, Dimetane)




Terfenadine (Seldane)

Neurologically-Active Drugs


- They affect the mechanism of message transfer from nerve to receptor.
- Example: Tranquilisers and analgesics

Tranquilisers

- Used for the treatment of stress and mental diseases
- Anti-depressant drugs are used when a person is suffering from depression. Example: Iproniazid, phenelzine




Iproniazid




Phenelzine (Nardil)


- Chlordiazepoxide and meprobamate are suitable for relieving tension, and equanil is used for controlling depression and hypertension.



Chlordiazepoxide



Meprobamate



Equanil

- Barbiturates (derivatives of barbituric acid viz., veronal, amytal, nembutal, luminal) constitute an important class of tranquilisers. They are hypotonic, i.e., sleep-producing agents.


Analgesics

- Used for reducing pain, without causing impairment of consciousness, mental confusion, incoordination or paralysis, or some other disturbances of the nervous system
- Two types: Non-narcotic (non-addictive) analgesics and narcotic drugs
- **Non-narcotic drugs**
-
- Example: Aspirin and paracetamol
- Aspirin inhibits the synthesis of chemicals known as prostaglandins, which stimulate inflammation in the tissue and cause pain.
- Aspirin also finds use in the prevention of heart attacks.

Reason: Because of its anti-blood-clotting action

-

- These drugs also reduce fever (**anti-pyretic**), and relieve skeletal pain occurring due to arthritis.
- **Narcotic drugs**
- Relieve pain and produce sleep (in medicinal doses)
- Produce stupor, coma, convulsions and ultimately death (in poisonous doses)
- Used for the relief of post-operative pain, cardiac pain and pains of terminal cancer, and in child birth
- Example: Morphine and its homologous such as heroin, codeine




Antimicrobials and Antifertility Drugs

Antimicrobials

- Inhibit the pathogenic action of microbes such as bacteria, fungi, virus and other parasites.
- Example: Antibiotics, antiseptics, disinfectants

Antibiotics


- Used for treating infections because of their low toxicity for humans and animals
- Inhibit the growth, and even destroy, microorganisms
- Sulpha drugs contain sulphanilamide, which is the real active compound. One of the most effective one is sulphapyridine.
- Structures of some antibiotics are as follows:



- Have either cidal (killing) effect or a static (inhibitory) effect on microbes. A few examples are listed in the given table.


Bactericidal	Bacteriostatic
Penicillin	Erythromycin
Aminoglycosides	Tetracycline
Ofloxacin	Chloramphenicol

- Antibiotics which kill or inhibit a wide range of gram-positive and gram-negative bacteria are called **broad-spectrum antibiotics**. Ampicillin and amoxicillin, which are synthetic modifications of penicillin, are broad-spectrum antibiotics.



General Structure of Penicillin

- Chloramphenicol is a broad-spectrum antibiotic which is used orally for the treatment of typhoid, dysentery, acute fever and pneumonia.



- Vancomycin and ofloxacin are broad-spectrum antibiotics.
- Dysidazirine, an antibiotic, is toxic towards certain strains of cancer cells.
- Antibiotics which are effective mainly against gram-positive or gram-negative bacteria are called narrow-spectrum antibiotics. Example: Penicillin G.
- Antibiotics which are effective against a single organism or disease are called limited-spectrum antibiotics.

Antiseptic

- Chemicals which either kill or prevent the growth of microorganisms
- Antiseptics are applied to living tissues such as wounds, cuts, ulcers and diseased skin surfaces.
- Example:
- Furacine
- Soframicine
- Dettol (mixture of chloroxylenol and terpineol)
- Tincture of iodine (2 – 3% solution of iodine in alcohol–water mixture): Applied on wounds
- Iodoform: An antiseptic for wounds
- Boric acid in dilute aqueous solution: Weak antiseptic for the eyes


Disinfectant

- Chemicals which either kill or prevent the growth of microorganisms
- Applied to inanimate objects such as floors, drainage system, instruments, etc.
- Example:
- 0.2 to 0.4 ppm of chlorine in aqueous solution
- Sulphur dioxide in low concentration


- 0.2% solution of phenol is used as an antiseptic while 1% solution of phenol is used as a disinfectant.

Antifertility Drugs

- Used in family planning
- Example: norethindrone, ethynylestradiol (novestrol)
- Norethindrone: Synthetic progesterone



- Ethynylestradiol (novestrol): Oestrogen derivative, which is used in combination with progesterone derivative



Chemicals in Food & Cleansing Agents

Chemicals in Food

Chemicals are added to food as –

- Food colours
- Flavours and sweeteners
- Fat emulsifiers and stabilising agents
- Flour improvers: anti-staling agents and bleaches

- Anti-oxidants
- Preservatives
- Nutritional supplements (like minerals, vitamins, amino acids)
- Artificial sweetening agents
- Chemicals that sweeten food
- Do not add calories to our body
- Saccharin (ortho-sulphobenzimide) is the first popular artificial sweetening agent.
- Some other artificial sweetening agents are aspartame, alitame, sucrolose, etc.
- Food preservatives
- Chemicals that prevent food from spoilage due to microbial growth
- Example: Table salt, vegetable oil, sodium benzoate (C_6H_5COONa), salts of propanoic acid


Antioxidants

Antioxidants are those compounds in food which prevent the damaging effect of free radicals in our body. They neutralise free radicals that are formed due to oxidation takes place in human body.

Free radicals are quite harmful and linked with diseases such as cancer and heart and liver diseases. Free radicals also damage the cells which lead to aging. Plant food that contains vitamin A and C such as fruits, vegetables, nuts are rich sources of antioxidants.


Cleansing Agents

- **Soaps**
- Soaps are sodium or potassium salts of long-chain fatty acids (such as stearic acid, oleic acid, palmitic acid).
- Saponification: Heating of fat (glyceryl ester of fatty acid) with aqueous solution of NaOH to form soap containing sodium salts



- Potassium soaps are soft to the skin than sodium soaps.
- Soaps do not work in hard water.



Reason: Hard water contains calcium and magnesium ions. When soaps are dissolved in hard water, these ions displace sodium or potassium from their salts and form insoluble calcium or magnesium salts of fatty acids. These insoluble salts separate as scum.




- **Synthetic detergents:** Work in both hard and soft water.

Classified into the following three categories –

- **Anionic detergents:** Sodium salts of sulphonated long-chain alcohols or hydrocarbons. Anionic part is involved in the cleansing action.
- Example:





- Cationic detergents: Quaternary ammonium salts of amines, with acetates, chlorides or bromides as anions.
- Example:



Cetyltrimethylammonium bromide


- Non-ionic detergents: Do not contain any ion.
- Example:




- The more the branching of chain in detergents, the more is the difficulty in their degradation.

Cleansing Action of Soap

Soap molecule has two ends- one polar or ionic end (the end with sodium or potassium ion) and one non-polar or non ionic end (the end with long hydrocarbon chain). The polar end is hydrophilic in nature i.e., this end is attracted towards water. The non-polar end is hydrophobic, but lipophilic in nature and it is attracted towards dirt.



When a soap molecule is dissolved in water, its hydrophobic ends attach themselves to dirt and remove it from the cloth, as shown sequentially in the figure that follows.



First, the non ionic part gets attached with dirt molecule and ionic part remains outside. In this way the molecules of soap arrange themselves in micelle formation and trap the dirt in the centre of the cluster. These micelles remain suspended in water like particles in a colloidal solution. Thus, the dust particles remain trapped in micelles (which remain suspended) and are easily rinsed away with water. Hence, soap micelles remove dirt by dissolving it in water.