

Question 1:

(a) Fill in the blanks by choosing the appropriate word/words from those given in the brackets: [4 x 1]

(iodoform, volume, mass, haloform, gram equivalent, chloroform, carbylmine, sp^3d^2 , high, coke, d^2sp^3 , low, gram mole, carbon monoxide.

(i) Equivalent conductivity is the conducting power of all the ions furnished by one _____ of an electrolyte present in a definite _____ of the solution.

(ii) Bleaching powder, on treatment with ethanol or acetone gives _____.

(iii) Outer orbital complexes involve _____ hybridization and are _____ spin complexes.

(iv) Zinc Oxide is reduced by _____ at 1673K to form zinc and _____.

(b) Select the correct alternative from the choices given: [4 x 1]

(i) The packing efficiency of simple cubic structure, body centered cubic structure and face centered cubic structure respectively is:

(1) 52.4%, 74%, 68%

(2) 74%, 68%, 52.4%

(3) 52.4%, 68%, 74%

(4) 68%, 74%, 52.4%

(ii) When acetone is treated with Grignard's reagent, followed by hydrolysis, the product formed is:

- (1) Secondary alcohol
- (2) Tertiary alcohol
- (3) Primary alcohol
- (4) Aldehyde

(iii) Which of the following electrolytes is least effective in causing flocculation of positively charged ferric hydroxidesol?

- (1) $K_3[Fe(CN)_6]$
- (2) K_2CrO_4
- (3) $K_4[Fe(CN)_6]$
- (4) KBr

(iv) On heating an aliphatic primary amine with chloroform and alcoholic potassium hydroxide, the organic compound formed is an:

- (1) Alkyl isocyanide
- (2) Alkanol
- (3) Alkanal
- (4) Alkyl cyanide

(c) Match the following:

[4 x 1]

- | | |
|-----------------------------|------------------------------|
| (i) Silicon and phosphorous | (a) Acetaldehyde |
| (ii) Iodoform test | (b) Xenon hexafluoride |
| (iii) Arrhenius equation | (c) n-type of semiconductors |

(iv) Distorted octahedral structure (d) Frequency factor

(d) Answer the following questions:

[4 x 2]

(i) What is the common name of the polymer obtained by the polymerization of caprolactam? Is it addition polymer or condensation polymer?

(ii) Why Zn^{2+} ions are colourless while Ni^{2+} ions are green and Cu^{2+} ions are blue in colour?

(iii) The molar conductivity of NaCl, CH_3COONa and HCl at infinite dilution is 126.45, 91.0 and 426.16 $ohm^{-1} cm^2 mol^{-1}$ respectively. Calculate the molar conductivity (λ_m^∞) for CH_3COOH at infinite dilution.

(iv) Identify the compounds A, B, C and D.

Question 2:

[2]

(a) An element has atomic weight $93 g mol^{-1}$ and density $11.5 g cm^{-3}$. If the edge length of its unit cell is 300 pm, identify the type of unit cell. ($N_A = 6.023 \times 10^{23} mol^{-1}$)

OR

(b) Calculate the radius of copper atom. The atomic weight of copper is $63.55 g mol^{-1}$. It crystallises in face centered cubic lattice and has density of $8.93 g cm^{-3}$ at 298. ($N_A = 6.023 \times 10^{23} mol^{-1}$)

Question 3:

[2]

Complete and balance the following chemical equations:



Question 4: [2]

(i) Write the chemical equation for the reaction of glucose with bromine water.

(ii) Write the zwitter ion structure of glucine.

Question 5: [2]

(i) How do antiseptics differ from disinfectants?

(ii) Name a substance that can be used as an antiseptic as well as a disinfectant.

Question 6: [2]

An alloy of gold (Au) and cadmium (Cd) crystallises with a cubic structure in which gold atoms occupy the corners and cadmium atoms fit into the face centres. What is the formula of this alloy?

Question 7: [2]

(a) State reasons for the following:

(i) Ethylamine is soluble in water whereas aniline is insoluble in water.

(ii) Aliphatic amines are stronger bases than aromatic amines.

OR

(b) Complete and balance the following equations:



Question 8: [2]

(a) Draw the structure of xenon tetrafluoride molecule. State the hybridisation of the central atom and the geometry of the molecule.

Question 9: [2]

(a) Calculate the emf and ΔG for the given cell at 25^o C:



$$\text{Given: } E_{\text{Cr}^{3+}/\text{Cr}}^{\circ} = -0.74\text{V}, \quad E_{\text{Fe}^{2+}/\text{Fe}}^{\circ} = -0.44\text{V}$$

$$(\text{IF} = 96500\text{C}, \text{R} = 8.314\text{JK}^{-1}\text{mol}^{-1})$$

OR

(b) Calculate the degree of dissociation (α) of acetic acid, if its molar conductivity (Δm) is 39.05 S cm² mol⁻¹

(Given: $\lambda_{(\text{H}^{+})}^{\circ} = 349.6\text{ S cm}^2\text{ mol}^{-1}$ and $\lambda_{(\text{CH}_3\text{COO}^{-})}^{\circ} = 40.95\text{ S cm}^2\text{ mol}^{-1}$)

Question 10: [3]

Name an important ore of silver. How is silver extracted from its sulphide ore? Give balanced chemical equations involved in the extraction of pure silver.

Question 11: [3]

How will you convert the following:

- (i) Chlorobenzene to biphenyl
- (ii) Propene to 1-bromopropane
- (iii) Chlorobenzene to aniline

Question 12: [3]

Explain what is observed when:

- (i) A beam of light is passed through a colloidal solution.
- (ii) An electric current is passed through a colloidal solution.
- (iii) An electrolyte (AlCl_3) is added to a colloidal solution of arsenious sulphide (As_2S_3).

Question 13: [3]

(a) How will you convert the following: (Give balanced equation)

- (i) Benzoyl chloride to benzaldehyde.
- (ii) Methyl chloride to acetic acid.
- (iii) Acetic acid to methane.

OR

(b) A ketone A ($\text{C}_4\text{H}_8\text{O}$) which undergoes Iodoform reaction gives compound B on reduction. B on heating with conc. H_2SO_4 at 443K gives a compound C which forms ozonic D. D on hydrolysis with Zn dust gives only E. Identify the compounds A to E. Write the Iodoform reaction with compound A.

Question 14: [3]

A first order reaction is 50% completed in 30 minutes at 300K and in 10 minutes at 320K. Calculate the activation energy of the reaction. ($R=3.814 \text{ JK}^{-1}\text{mol}^{-1}$).

Question 15:**[3]**

Explain the following:

- (i) Transition metals and their compounds generally exhibit a paramagnetic behaviour.
- (ii) There is an increase in density of elements from titanium ($Z=22$) to copper ($Z=29$) in the 3d series of transition elements.
- (iii) $K_2Cr_2O_7$ acts as a powerful oxidising agent in acidic medium.

Question 16:**[3]**

(a) (i) The elevation in boiling point when 0.30 g of acetic acid is dissolved in 100 g of benzene is $0.0633^\circ C$. Calculate the molecular weight of acetic acid from this data. What conclusion can you draw about the molecular state of the solute in the solution?

(Given K_b for benzene = $2.53K \text{ kg mol}^{-1}$, at. Wt. of C=12, H=1, O=16)

(ii) Determine the osmotic pressure of a solution prepared by dissolving 0.025 g of K_2SO_4 in 2 litres of water at $25^\circ C$, assuming that K_2SO_4 is completely dissociated.

($R=0.0821 \text{ Lit-atm K}^{-1} \text{ mol}^{-1}$, mol. wt. of $K_2SO_4 = 174 \text{ g mol}^{-1}$)

OR

(b) (i) An aqueous solution of a non-volatile solute freezes at 272.4 K , while pure water freezes at 273.0 K . Determine the following:

(Given $K_f = 1.86 \text{ K kg mol}^{-1}$, $K_b = 0.512 \text{ K kg mol}^{-1}$ and vapour pressure of water at $298 \text{ K} = 23.756 \text{ mm of Hg}$)

- (1) The molality of solution
- (2) Boiling point of solution
- (3) The lowering of vapour pressure of water at 298 K

(ii) A solution containing 1.23 g of calcium nitrate in 10 g of water, boils at 100.975°C at 760 mm of Hg . Calculate the van't Hoff factor for the salt at this concentration.

(K_b for water = $0.52 \text{ K kg mol}^{-1}$, mol. wt. of calcium nitrate = 164 g mol^{-1})

Question 17:

[3]

(a) (i) Write the IUPAC names of the following complexes:



(ii) With reference to the coordination complex ion $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$ answer the following: (at. No. of $\text{Fe} = 26$)

- (1) Give the IUPAC name of the complex ion.
- (2) What is the oxidation number of the central metal ion?
- (3) How many unpaired electrons are there in the complex ion?
- (4) State The type of hybridisation of the complex ion.

OR

(b) (i) Name of the type of isomerism exhibited by the following pairs of compounds:

- (1) $[\text{Co}(\text{ONO})(\text{NH}_3)_5]^{2+}$ and $[\text{Co}(\text{NO}_2)(\text{NH}_3)_5]^{2+}$
- (2) $[\text{Cr}(\text{H}_2\text{O})_4\text{Cl}_2] \cdot \text{Cl} \cdot 2\text{H}_2\text{O}$ and $[\text{Cr}(\text{H}_2\text{O})_5\text{Cl}]\text{Cl}_2 \cdot \text{H}_2\text{O}$
- (3) $[\text{Co}(\text{NH}_3)_6][\text{Cr}(\text{CN})_6]$ and $[\text{Cr}(\text{NH}_3)_6][\text{Co}(\text{CN})_6]$

(ii) Using the valence bond approach, predict the shape, hybridisation and magnetic behaviour of $[\text{Ni}(\text{CO})_4]$. (at. no. of Ni=28)

Question 18:

[3]

(a) (i) Give balanced chemical equations for the following reactions:

- (1) Phenol is treated with ice cold alkaline solution of benzene diazonium chloride.
- (2) Diethyl ether is treated with phosphorous pentachloride.
- (3) Ethyl alcohol is treated with thionyl chloride.

(ii) Give one chemical test each to distinguish between the following pairs of compounds:


- (1) Ethanol and dimethyl ether
- (2) Propan-1-ol and propan-2-ol

OR

(b) (i) Write chemical equations to illustrate the following name reactions:

- (1) Williamson's synthesis
- (2) Esterification reaction
- (3) Reimer-Tiemann reaction

(ii) Identify the compounds A and B in the given reactions:



www.ncertbooks.net