

ISC Paper 2017

Chemistry

Maximum Marks: 70

Time allowed: 3 hours

- Answer all questions in Part I and six questions from Part II, choosing two questions from Section A, two from Section B and two from Section C.
 - All working, including rough work, should be done on the same sheet as, and adjacent to, the rest of the answer.
 - The intended marks for questions or parts of questions are given in brackets [].
 - Balanced equations must be given wherever possible and diagrams where they are helpful.
 - When solving numerical problems, all essential working must be shown.
 - In working out problems use the following data:
Gas constant $R = 1.987 \text{ cal deg}^{-1} \text{ mol}^{-1} = 8.314 \text{ JK}^{-1} \text{ mol}^{-1} = 0.0821 \text{ dm}^3 \text{ atm K}^{-1} \text{ mol}^{-1}$. $1 \text{ L atm} = 1 \text{ dm}^3 \text{ atm} = 101.3 \text{ J}$. $1 \text{ Faraday} = 96500 \text{ Coulombs}$.
Avogadro's number = 6.023×10^{23}
-

Part – I (20 Marks)

Answer all questions.

Question 1.

(a) Fill in the blanks by choosing the appropriate word/words from those given in the brackets: [5]

(iodoform, acetaldehyde, positive, greater, acidic, acetone, disaccharide, negative, increases, glucose, decreases, chloroform, polysaccharide, lactose, lesser, basic, cationic hydrolysis, anionic hydrolysis)

(i) Calcium acetate on heating gives which gives on heating with iodine and sodium hydroxide solution.

(ii) On dilution of a solution, its specific conductance while its equivalent conductance

(iii) Sucrose is a and yields upon hydrolysis, a mixture of and fructose.

(iv) More the standard reduction potential of a substance, the is its ability to displace hydrogen from acids.

(v) An aqueous solution of CH_3COONa is due to

(b) Complete the following statements by selecting the correct alternative from the choices given: [5]

(i) In a face-centred cubic lattice, atom (A) occupies the corner positions and atom (B)

occupies the face centre positions. If one atom of (B) is missing from one of the face-centred points, the formula of the compound is:

- (1) A_2B_5
- (2) A_2B_3
- (3) AB_2
- (4) A_2B

(ii) The half-life period of a first-order reaction is 20 minutes. The time required for the concentration of the reactant to change from 0.16 M to 0.02 M is:

- (1) 80 minutes
- (2) 60 minutes
- (3) 40 minutes
- (4) 20 minutes

(iii) For a spontaneous reaction ΔG° and E° cell will be respectively:

- (1) -ve and +ve
- (2) +ve and -ve
- (3) +ve and +ve
- (4) -ve and -ve

(iv) The conjugate acid of HPO_4^{2-} is:

- (1) H_3PO_3
- (2) H_3PO_4
- (3) $H_2PO_4^-$
- (4) PO_4^{3-}

(v) The polymer formed by the condensation of hexamethylenediamine and adipic acid is:

- (1) Teflon
- (2) Bakelite
- (3) Dacron
- (4) Nylon-66

(c) Answer the following questions: [5]

(i) Why the freezing point depression (ΔT_f) of 0.4 M NaCl solution is nearly twice than that of 0.4 M glucose solution?

(ii) Identify the order of reaction from each of the following units of rate constant (k):

- (a) $\text{mol L}^{-1} \text{sec}^{-1}$
- (b) $\text{mol}^{-1} \text{L sec}^{-1}$

(iii) Specific conductivity of 0.20 M solution of KCl at 298 K is 0.025 S cm^{-1} . Calculate its molar conductivity.

(iv) Name the order of reaction which proceeds with a uniform rate throughout.

(v) What are the products formed when phenol and nitrobenzene are treated separately with a mixture of concentrated sulphuric acid and concentrated nitric acid?

(d) Match the following: [5]

(i) Diazotisation	(a) Bakelite
(ii) Argentite	(b) Nernst equation
(iii) Thermosetting plastics	(c) Aniline
(iv) Electrochemical cell	(d) Ethylenediamine
(v) Bidentate ligand	(e) Froth floatation process

Answers:

- (a) (i) acetone, iodoform
(ii) decreases, increases
(iii) disaccharide, glucose
(iv) negative, greater
(v) basic, anionic hydrolysis

- (b) (i) (1)
(ii) (2)
(iii) (1)
(iv) (3)
(v) (4)

(c) (i) This is because Van't Hoff factor for NaCl is 2 and for glucose, it is 1.

(ii) (a) – Zero

(b) – Two

(iv) Zero

(v) 2, 4, 6 – Trinitrophenol

m – Dinitrobenzene

(d) (i) – (c)

(ii) – (e)

(iii) – (a)

(iv) – (b)

(v) – (d)

Part – II (50 Marks)

Section – A

Answer any two questions.

Question 2.

(a) (i) Determine the freezing point of a solution containing 0.625 g of glucose ($C_6H_{12}O_6$) dissolved in 102.8 g of water. [2]

(Freezing point of water = 273 K, K_f for water = $1.87 \text{ K kg mol}^{-1}$, at. wt. C = 12, H = 1, O = 16)

(ii) A 0.15 M aqueous solution of KCl exerts an osmotic pressure of 6.8 atm at 310 K. [2] Calculate the degree of dissociation of KCl. ($R = 0.0821 \text{ Lit. atm K}^{-1} \text{ mol}^{-1}$).

(iii) A solution containing 8.44 g of sucrose in 100 g of water has a vapour pressure 4.56 mm of Hg at 273 K. If the vapour pressure of pure water is 4.58 mm of Hg at the same temperature, calculate the molecular weight of sucrose. [1]

(b) (i) When ammonium chloride and ammonium hydroxide are added to a solution containing both Al^{3+} and Ca^{2+} ions, which ion is precipitated first and why? [2]

(ii) A solution of potassium chloride has no effect on litmus whereas, a solution of zinc chloride turns the blue litmus red. Give a reason. [2]

(c) How many sodium ions and chloride ions are present in a unit cell of sodium chloride crystal? [1]

Answer:

(a) (i) $M_B = 180 \text{ U}$

$$\Delta T_f = K_f \cdot \frac{W_B \times 1000}{M_B \times W_A}$$

$$= \frac{1.87 \times 0.625 \times 1000}{180 \times 102.8} = \frac{1.87 \times 625}{180 \times 102.8} \text{ K} = 0.06 \text{ K}$$

$$T_{f'} = 273 - 0.06 = 272.94 \text{ K}$$

(ii) $R = 0.0821 \text{ L atm. K}^{-1} \text{ mol}^{-1}$

$$\pi = i CRT$$

$$6.8 = i \times 0.15 \times 0.0821 \times 310$$

$$i = 1.78$$

$$\alpha = \frac{i-1}{n-1}$$

For KCl, $n = 2$

$$\alpha = \frac{1.78-1}{2-1}$$

Degree of dissociation, $\alpha = 0.78$ or 78%

(iii) $P_A^\circ = 4.58 \text{ mm of Hg}$
 $P_A = 4.56 \text{ mm of Hg}$

$$\frac{P_A^\circ - P_A}{P_A^\circ} = \chi_B = \frac{n_B}{n_B + n_A}$$

$$\frac{4.58 - 4.56}{4.58} = \frac{\frac{8.44}{M_B}}{\frac{100}{18} + \frac{8.44}{M_B}}$$

$$\frac{0.02}{4.58} = \frac{8.44}{M_B} \times \frac{18M_B}{(100M_B + 151.92)}$$

$$\frac{0.02}{4.58} = \frac{8.44 \times 18}{100M_B + 151.92}$$

$$100M_B + 151.92 = \frac{8.44 \times 18 \times 4.58}{0.02}$$

$$M_B = 346.38 \text{ amu}$$

(b) (i) Al^{3+} ions are precipitated first as $\text{Al}(\text{OH})_3$ because K_{sp} for $\text{Al}(\text{OH})_3$ is lower than that of $\text{Ca}(\text{OH})_2$.

(ii) This is because KCl does not undergo hydrolysis in water and the aqueous solution is neutral whereas ZnCl_2 undergoes hydrolysis in water to give an acidic solution which turns blue litmus solution red.

(c) Number of Na^+ ions = 4
 Number of Cl^- ions = 4

Question 3.

(a) (i) Lead sulphide has a face-centred cubic crystal structure. If the edge length of the unit cell of lead sulphide is 495 pm, calculate the density of the crystal. [1]

(at. wt. of Pb = 207, S = 32)

(ii) For the reaction: $2\text{H}_2 + 2\text{NO} \rightleftharpoons 2\text{H}_2\text{O} + \text{N}_2$, the following rate data was obtained: [3]

S.No.	[NO] mol L ⁻¹	[H ₂] mol L ⁻¹	Rate : mol L ⁻¹ sec ⁻¹
1	0.40	0.40	4.6×10^{-3}
2	0.80	0.40	18.4×10^{-3}
3	0.40	0.80	9.2×10^{-3}

Calculate the following:

(1) The overall order of a reaction.

- (2) The rate law.
 (3) The value of rate constant (k).
 (b) (i) The following electrochemical cell is set up at 298 K: [2]

- (1) Write the cell reaction.
 (2) Calculate the emf and free energy change at 298 K.
 (a) Answer the following: [2]
 (1) What is the effect of temperature on the ionic product of water (K_w)?
 (2) What happens to the ionic product of water (K_w) if some acid is added to it?
 (c) Frenkel defect does not change the density of the ionic crystal whereas, Schottky defect lowers the density of ionic crystal. Give a reason; [2]

Answer:

(a) (i)
$$\rho = \frac{Z \times M}{N_0 a^3}$$

$$a = 495 \times 10^{-10} \text{ cm}$$

$$M = 207 + 32 = 239 \text{ U}$$

$$N_0 = 6.022 \times 10^{23}$$

$$Z = 4$$

$$\rho = \frac{4 \times 239}{6.022 \times 10^{23} \times (4.95)^3 \times 10^{-24}}$$

$$= \frac{4 \times 2390}{6.022 \times (4.95)^3} = \frac{4 \times 2390}{6.022 \times 121.29} = \frac{7560}{730.41}$$

$$\rho = 10.35 \text{ g/cc}$$

(ii) Let rate law is

$$r = k[\text{H}_2]^x [\text{NO}]^y$$

$$4.6 \times 10^{-3} = k (0.4)^x (0.4)^y \quad \dots(1)$$

$$18.4 \times 10^{-3} = k (0.4)^x (0.80)^y \quad \dots(2)$$

$$9.2 \times 10^{-3} = k (0.8)^x (0.4)^y \quad \dots(3)$$

Dividing equation (1) by equation (2), we have

$$\frac{1}{4} = \left(\frac{1}{2}\right)^y$$

$$y = 2$$

Dividing equation (1) by equation (3), we have

$$\frac{1}{2} = \left(\frac{1}{2}\right)^x$$

$$x = 1$$

Therefore,

Order w.r.t. $H_2 = 1$

Order w.r.t. $NO = 2$

Overall order = $1 + 2 = 3$

Rate law $r = k[H_2]^1 [NO]^2$



From equation (1)

$$4.6 \times 10^{-3} = k(0.4)^1 (0.4)^2$$

$$4.6 \times 10^{-3} = k \times 4 \times 4 \times 4 \times 10^{-3}$$

$$k = \frac{4.6}{64}$$

$$k = 0.07 \text{ mol}^{-2} \text{ L}^2 \text{ sec}^{-1}$$



(2) $E_{\text{cell}}^{\circ} = 0.339 + 0.761$

$$= 1.10 \text{ V}$$

$$\Delta G^{\circ} = -n F E_{\text{cell}}^{\circ}$$

$$= -2 \times 96500 \times 1.10$$

$$\Delta G^{\circ} = -212.3 \text{ kJ}$$

(ii) (1) K_w for water increases with the increase in temperature due to the increase in the degree of ionisation of water.

(2) K_w remains unchanged.

(c) This is because in Frenkel defect no ions are missing from the crystal lattice site whereas, in case of Schottky defect, an equal number of positive and negative ions are missing from the crystal lattice, hence density decreases.

Question 4.

- (a) (i) Name the law or principle to which the following observations confirm: [3]
- (1) When water is added to a 1.0 M aqueous solution of acetic acid, the number of hydrogen ion (H^+) increases.
- (2) When 9650 coulombs of electricity is passed through a solution of copper sulphate, 3.175 g of copper is deposited on the cathode (at. wt. of Cu = 63.5).
- (3) When ammonium chloride is added to a solution of ammonium hydroxide, the concentration of hydroxyl ion decreases.
- (ii) What is the difference between the order of a reaction and its molecularity? [2]
- (b) (i) Explain why high pressure is required in the manufacture of sulphur trioxide by the contact process. State the law or principle used. [2]
- (ii) Calculate the equilibrium constant (K_c) for the formation of NH_3 in the following reaction: [1]
- $$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$$
- At equilibrium, the concentration of NH_3 , H_2 and N_2 are 1.2×10^{-2} , 3.0×10^{-2} and 1.5×10^{-2} M respectively.
- (c) Explain the following: [2]
- (i) Hydrolysis of ester (ethyl acetate) begins slowly but becomes fast after some time.
- (ii) The pH value of acetic acid increases on the addition of a few drops of sodium acetate.

Answers:


(a) (i) (1) This is due to the increase in the degree of ionisation of CH_3COOH .

$$(2) W = ZQ \\ = \frac{63.5}{2 \times 96500} \times 9650 \\ W = 3.175 \text{ g}$$

(3) This is due to the common ion effect (NH_4^+ ions are common)

(ii)

Order of a reaction	Molecularity
<ol style="list-style-type: none">1. It can be zero.2. It can be the whole number or fractional.3. It is defined for simple as well as complex reaction.4. It is determined experimentally.	<ol style="list-style-type: none">1. It is never zero.2. It is always a whole number.3. It is defined only for simple reactions.4. It is a theoretical concept.




This is because the forward reaction is accompanied by a decrease in volume.

The principle used is i.e., Chatelier's principle.

It states that when a system in equilibrium is subjected to stress, (i.e., change of

concentration, temperature pressure etc.), the equilibrium tends to shift in a direction so as to undergo the effect of applied stress.



$$\begin{aligned}K_c &= \frac{[\text{NH}_3]^2}{[\text{N}_2][\text{H}_2]^3} = \frac{(1.2 \times 10^{-2})^2}{(1.5 \times 10^{-2})(3 \times 10^{-2})^3} \\&= \frac{1.2 \times 1.2 \times 10^{-4}}{1.5 \times 10^{-2} \times 3 \times 3 \times 3 \times 10^{-6}} = \frac{1.2 \times 1.2 \times 10^4}{1.5 \times 9 \times 3} \\&= \frac{1.44 \times 10^4}{1.5 \times 9 \times 3} = 400\end{aligned}$$

(c) (i) This is because of H^+ ions produced during hydrolysis act as a catalyst.

(ii) This is due to the common ion effect. As a result of (H^+) decreases.

Section – B

Answer any two questions.

Question 5.

(a) Write the formula of the following compounds: [2]

(i) Potassium trioxalatoaluminate (III).

(ii) Hexaaquairon (II) sulphate.

(b) Name the types of isomerism shown by the following pairs of compounds: [1]

(i) $[\text{Cu}(\text{NH}_3)_4][\text{PtCl}_4]$ and $[\text{Pt}(\text{NH}_3)_4][\text{CuCl}_4]$

(ii) $[\text{Co}(\text{Pn})_2\text{Cl}_2]^+$ and $[\text{Co}(\text{en})_2\text{Cl}_2]^+$

(c) For the coordination complex ion $[\text{Co}(\text{NH}_3)_6]^{3+}$ [2]

(i) Give the IUPAC name of the complexion.

(ii) What is the oxidation number of cobalt in the complexion?

(iii) State the type of hybridisation of the complexion.

(iv) State the magnetic behaviour of the complexion.

Answers:

(a) (i) $\text{K}_3[\text{Al}(\text{C}_2\text{O}_4)_3]$

(ii) $[\text{Fe}(\text{H}_2\text{O})_6] \text{SO}_4$

(b) (i) Coordination isomerism

(ii) Linkage isomerism

(c) (i) hexaamminecobalt (II) ions

(ii) $x + 6(0) = +3$

$x = +3$


No. of Co in complex ion = +3

- (iii) d^2sp^3
 (iv) $[\text{Co}(\text{NH}_3)_6]^{3+}$
 Co^{3+} has no unpaired electron.
 Hence, $[\text{Co}(\text{NH}_3)_2]^{3+}$ is diamagnetic.

Question 6.


- (a) Give balanced equations for the following reactions: [3]
 (i) Potassium permanganate is heated with concentrated hydrochloric acid.
 (ii) Lead sulphide is heated with hydrogen peroxide.
 (iii) Ozone is treated with potassium iodide solution.
 (b) Discuss the theory involved in the manufacture of sulphuric acid by the contact process. [2]

Answers:




(b) **Theory of Contact process :**


(i) **Production of SO_2 :**



(ii) **Oxidation of SO_2 to SO_3 :**






(iii) **Conversion of SO_3 into H_2SO_4 :**




Question 7.

- (a) (i) What are the types of hybridisation of iodine in interhalogen compounds IF_3 , IF_5 and IF_7 , respectively? [3]
 (ii) Draw the structure of xenon hexafluoride (XeF_6) molecule and state the hybridisation of the central atom.
 (b) Give the balanced equations for the conversion of argentite (Ag_2S) to metallic silver. [2]



Answers:



Distorted octahedral geometry.



Argentite



Section - C

Answer any two questions.

Question 8.

(a) How can the following conversions be brought about:

(i) Acetaldehyde to propan-2-ol. [1]

(ii) Nitrobenzene to p-aminoazobenzene. [1]


(iii) Acetic acid to methylamine. [2]

(iv) Aniline to benzene. [1]






(b) (i) How will you distinguish between primary, secondary and tertiary amines by Hinsberg's test? [1]

(ii) Why do alcohols possess higher boiling points as compared to those of corresponding alkanes? [1]

(c) Identify the compounds A, B and C: [3]



Answer:




(b) (i) With Hinsberg's reagent:

Primary amines give N-alkyl benzene sulphonamide soluble in alkali.

Secondary amines give N, N-dialkyl benzene sulphonamide insoluble in alkali.


Tertiary amines have no action with Hinsberg's reagent.

(ii) This is because in alcohols there are intermolecular H-bonds which are stronger than Van der Waals forces of attraction in alkanes.



(c) (i) A $\text{C}_6\text{H}_5\text{COCl}$, Benzoyl chloride

B $\text{C}_6\text{H}_5\text{CHO}$, Benzaldehyde



(ii) A — CH_3CHO , Acetaldehyde

B — CH_3CH_2 , Ethyl alcohol

OH

C — $\text{C}_2\text{H}_5 - \text{O} - \text{C}_2\text{H}_5$, Diethyl ether

Question 9.

(a) Give balanced equations for the following name reactions: [3]

(i) Friedel-Crafts reaction (alkylation)

(ii) Williamson's synthesis

(iii) Aldol condensation

(b) Give the chemical test to distinguish: [3]

(i) Ethyl alcohol and sec-propyl alcohol

(ii) Acetaldehyde and acetic acid

(c) (i) Deficiency of which vitamin causes the following diseases: [4]

(1) Scurvy

(2) Night blindness

(ii) Write two differences between globular and fibrous proteins.

(a) Name the two organic compounds which have the same molecular formula C_2H_6O . Will they react with PCl_5 ? If they react, what are the products formed?

(c) Give balanced equations for the following reactions: [3]

(i) Methyl magnesium bromide with ethyl alcohol.

(ii) Acetic anhydride with phosphorus pentachloride.

(iii) Acetaldehyde with hydroxylamine.

Answer:

(a) (i) A – $CH_3 - C \equiv CH$, propyne


B – CH_3COCH_3 , Acetone

C – CH_3COOH , Acetic acid

D – $HCOOH$, Formic acid (or $CO_2 + H_2O$)

E – CH_3COCl , Acetyl chloride


F – $CH_3COOC_2H_5$, Ethyl acetate



The above reaction is Hell - Volhard - Zelinsky.

(b) (i) Nylon - 6, Condensation polymer.

(ii) C_2H_5OH , Ethyl alcohol and CH_3OCH_3 Dimethyl ether. C_2H_5OH gives C_2H_5Cl with PCl_5 .



CH_3OCH_3 gives CH_3Cl with PCl_5 .

