
Study of Acids, Bases and Salts

IMPORTANT POINTS TO REMEMBER

- 1. Acids: The chemical compounds which on dissolving in water produce Hydrogen ions or Hydronium ions as the only positively charged particles. All acids essentially contain Hydrogen.
- 2. Hydronium ion: Hydronium ion is hydrated hydrogen ion. When hydrogen ion combines with water molecule it forms hydrated hydrogen ion called hydronium ion. The formation of hydronium ion takes place by the coordinate bond formation. In water molecule, oxygen atom has two lone pairs of electrons, oxygen atom shares one of its lone pair of electrons to hydrogen ion to complete its dublet thereby the formation of hydronium ion takes place.
 - (i) Formation of Water molecule



(ii) Formation of Hydrogen ion (H⁺)

$$H - e^- \longrightarrow H^+$$

(iii) Formation of Hydronium ion

$$H_2O + H^+ \longrightarrow H_3O^+$$
 $H \stackrel{\circ}{\longrightarrow} H$
 $H \stackrel{\circ}{\longrightarrow} H$
 H
 H
 H
 H
 H

- 3. Acids can be classified on the basis of
 - (i) Strength of an acid (ii) Basicity of an acid (iii) Concentration
- 4. Strong acids are those which almost completely ionized in their aqueous solutions. The solution of strong acid contains mostly ions. e.g., Hydrochloric acid (HCl), Sulphuric acid (H2SO4), Nitric acid (HNO₃).
- 5. Weak acids are those which partially or incompletely ionized in their aqueous solutions. The solution of weak acid contains ions as well as molecules, e.g., Acetic acid (CH3COOH), Sulphurous acid (H2SO3), Carbonic acid (H₂CO₃).
- 6. Basicity of an acid is defined as the number of replaceable Hydrogen ions present per molecule of an acid in its aqueous solution.
- 7. Monobasic acids: Acids which contain only one replaceable Hydrogen ion per molecule are called Monobasic acids. e.g., HCl, HNO3, CH3COOH

Monobasic acids form single series of salts, i.e., normal salts

NaOH + HCl
$$\longrightarrow$$
 NaCl + H₂O
HCl \longrightarrow H⁺ + Cl⁻
NaOH + HNO₃ \longrightarrow NaNO₃ + H₂O
HNO₃ \longrightarrow H⁺ + NO₃⁻
NaOH + CH₃COOH \longrightarrow CH₃COONa + H₂O
CH₃COOH \longrightarrow CH₃COO⁻ + H⁺

8. The names of the salts formed by some monobasic acids are given below:

	Monobasic acids	Salts
(<i>i</i>)	Nitric acid	Nitrate
(ii)	Nitrous acid	Nitrite
(iii)	Hydrochloric acid	Chloride
(iv)	Acetic acid	Acetate
(v)	Hydrobromic acid	Bromide
(vi)	Hydrofluoric acid	Fluoride

9. Dibasic acids are those which contain two replaceable Hydrogen ions per molecule of an acid in its aqueous solution, e.g, H₂SO₄, H₂SO₃, H₂CO₃, etc.
For example (i) Ionization of Sulphuric acid

$$\begin{array}{c} H_2SO_4 \longrightarrow H^{\dagger} + HSO_4^{\dagger} \\ \hline HSO_4^{\dagger} \longrightarrow H^{\dagger} + SO_4^{2-} \\ \hline H_2SO_4 \longrightarrow 2H^{\dagger} + SO_4^{2-} \end{array}$$

Dibasic acids form two series of salts, one acid salt and one normal salt

$$NaOH + H_2SO_4 \longrightarrow NaHSO_4 + H_2O$$
Sodium bisulphate

(Acid salt)

 $NaHSO_4 + NaOH \longrightarrow Na_2SO_4 + H_2O$

Sodium sulphate

(Normal salt)

(ii) The ionization/dissociation of sulphurous acid

$$H_2SO_3 \Longrightarrow H^+ + HSO_3^ H_2SO_3 \Longrightarrow 2H^+ + SO_3^{2-}$$
 $H_2SO_3 \Longrightarrow 2H^+ + SO_3^{2-}$
 $NaOH + H_2SO_3 \longrightarrow NaHSO_3 + H_2O$
 $Sodium bisulphite$
 $NaHSO_3 + NaOH \longrightarrow Na_2SO_3 + H_2O$
 $Sodium sulphite$

10. The names of the salts formed by some dibasic acids are given below:

Dibasic acids	Salts
(i) Sulphuric acid	Bisulphate & Sulphate
(ii) Sulphurous acid	Bisulphite & Sulphite
(iii) Carbonic acid	Bicarbonate & Carbonate
(iv) Hydrogen sulphide	Bisulphide & Sulphide

11. Tribasic acids: The acids which ionize on dissolving in water to produce three replaceable Hydrogen ions per molecule of an acid are called tribasic acids, e.g., H₃PO₄.

Ionization of Phosphoric acid

$$H_3PO_4 \Longrightarrow H^+ + H_2PO_4^ H_2PO_4^- \Longrightarrow H^+ + PO_4^{2-}$$
 $HPO_4^{2-} \Longrightarrow H^+ + PO_4^{3-}$
 $H_3PO_4 \Longrightarrow 3H^+ + PO_4^{3-}$

Tribasic acids form two different acid salts and one normal salt.

$$\begin{array}{c} \text{NaOH} \ + \ \text{H}_3\text{PO}_4 \ \longrightarrow \ \text{NaH}_2\text{PO}_4 \ + \ \text{H}_2\text{O} \\ & \text{(Acid salt)} \end{array}$$

$$\begin{array}{c} \text{NaH}_2\text{PO}_4 + \text{NaOH} \longrightarrow \ \text{Na}_2\text{HPO}_4 \ + \ \text{H}_2\text{O} \\ & \text{(Acid salt)} \end{array}$$

$$\begin{array}{c} \text{Na}_2\text{HPO}_4 + \text{NaOH} \longrightarrow \ \text{Na}_3\text{PO}_4 \ + \ \text{H}_2\text{O} \\ & \text{(Normal salt)} \end{array}$$

- 12. Concentration of an acid is the measure of the amount of water present in the acid.
- 13. Concentrated acid contains negligible quantities of water.
- 14. Dilute acid contains more quantity of water and less quantity of acid.
- 15. Strong acid will remains strong even if the solution is diluted and weak acid will remains weak even if the solution is concentrated.
 - 16. Methods of preparation of acids: Acids are generally prepared by the following methods:
 - (a) By synthesis or direct combination:

(b) By dissolving acidic oxide or acid anhydride in water:

Acidic oxides are also called as acid anhydrides. Acidic oxides on dissolving in water produces acids.

(c) By the reaction of non-volatile acid with the salts of more volatile acids:

Concentrated sulphuric acid is the non-volatile or least volatile acid which reacts with metallic nitrates and metallic chlorides to form Nitric acid and Hydrochloric acid respectively.

NaCl +
$$H_2SO_4$$
(conc.) \xrightarrow{below} NaHSO₄ + HCl Hydrochloric acid

KNO₃ + H_2SO_4 (conc.) \xrightarrow{below} KHSO₄ + HNO₃ Nitric acid

(d) By the oxidation of non-metals: Non-metals on reaction with concentrated acids get oxidized to respective oxyacids.

$$C + 2H_2SO_4(conc.) \longrightarrow CO_2 + 2H_2O + 2SO_2$$
Sulphur dioxide
$$C + 4HNO_3(conc.) \longrightarrow CO_2 + 2H_2O + 4NO_2$$
Nitrogen dioxide
$$S + 6HNO_3(conc.) \longrightarrow H_2SO_4 + 2H_2O + 6NO_2$$
Sulphuric acid
$$P + 5HNO_3(conc.) \longrightarrow H_3PO_4 + 5NO_2 + H_2O$$
Phosphoric acid

17. The physical properties of an acid are

- (i) They are sour in taste.
- (ii) They are corrosive in nature.
- (iii) They show characteristic colour change with indicators.

Acids show characteristic colour change with indicators.

Name of Indicator	Colour change	
1. Blue litmus	Red	
2. Phenolphthalein	Colourless	
3. Methyl orange	Red or Pink	

18. The typical chemical properties of a dilute acid are :

(a) Reaction with active metals: Dilute acids (except dilute nitric acid) on reaction with active metals produce a colourless and odourless gas which burns with a popping sound.

Nitric acid on reaction with metals produce oxides of nitrogen and water as it is a strong oxidizing agent. Only Magnesium and Manganese react with very dilute nitric acid to liberate hydrogen.

$$Mg + 2HNO_3 (v dil.) \longrightarrow Mg(NO_3)_2 + H_2 \uparrow$$

 $Mn + 2HNO_3 (v dil.) \longrightarrow Mn(NO_3)_2 + H_2 \uparrow$

(b) Reaction with bases: Dilute acids react with bases to form salt and water as the only products and the reaction is called as neutralization reaction.

$$KOH + HNO_3(dil.) \longrightarrow KNO_3 + H_2O$$

$$NH_4OH + HNO_3(dil.) \longrightarrow NH_4NO_3 + H_2O$$

$$2NaOH + H_2SO_4(dil.) \longrightarrow Na_2SO_4 + 2H_2O$$

Ionic equation

$$Na^+OH^- + H^+Cl^- \longrightarrow Na^+Cl^- + H_2O$$

Deleting the spectator ions

$$OH^- + H^+ \longrightarrow H_2O$$

In neutralization reaction, Hydroxyl ion derived from base and Hydrogen ion derived from an acid combine to form unionized water molecule.

(c) Reaction with metallic oxides: Metallic oxides react with acids to form salt and water.

(d) Reaction with metallic carbonates and bicarbonates: Dilute acids react with metallic carbonates and bicarbonates to liberate a colourless, odourless gas with brisk effervescence which turns lime water milky.

(e) Reaction with metallic sulphites and bisulphites: Dilute acids react with metallic sulphites and bisulphites to liberate a colourless gas having burnt Sulphur smell which turns acidified Potassium dichromate from orange to green and Potassium permanganate solution from purple to colourless, i.e., it decolourizes.

(f) Reaction with metallic sulphides and bisulphides: Dilute acids combine with metallic sulphides and bisulphides to form a colourless gas having rotten egg smell which turns lead acetate solution black.

$$FeS + 2HCl(dil.) \longrightarrow FeCl_2 + H_2S$$

$$Ferrous Hydrogen$$

$$chloride sulphide$$

$$NaHS + HCl (dil.) \longrightarrow NaCl + H_2S$$

$$Sodium Hydrogen$$

$$bisulphide chloride sulphide$$

$$2KHS + H_2SO_4 (dil.) \longrightarrow K_2SO_4 + 2H_2S$$

$$Potassium Potassium Hydrogen$$

$$bisulphide sulphide$$

$$H_2S + (CH_3COO)_2Pb \longrightarrow 2CH_3COOH + PbS$$

$$Lead acetate (Colourless)$$

$$Lead sulphide (Black)$$

- 19. The metallic hydroxides and metallic oxides are called bases. The general formula of metallic hydroxide is M(OH), where x is the valency of metal.
- 20. Soluble bases are called alkalies, e.g., NaOH, KOH, NH4OH and Ca(OH)2. Ca(OH)2 is sparingly soluble.
- 21. Alkalies: These are the soluble bases. The compounds which on dissolving in water produce hydroxyl ions as the only negatively charged particles.

$$NaOH \longrightarrow Na^{+} + OH^{-}$$
 $KOH \longrightarrow K^{+} + OH^{-}$
 $NH_{4}OH \longrightarrow NH_{4}^{+} + OH^{-}$

- 22. Ammonium ion: The ammonium ion is formed between ammonia molecule and hydrogen ion. In ammonia there are three single covalent bonds between nitrogen and hydrogen and one lone pair of electrons on nitrogen atom. The hydrogen ion shares lone pair of electrons from nitrogen atom to complete its dublet by a coordinate bond formation thereby leading to the formation of an ammonium ion.
 - (i) Formation of Ammonia molecule

(ii) Formation of Hydrogen ion

$$H + e^- \longrightarrow H$$

(iii) Formation of Ammonium ion

$$\begin{array}{c} H \\ \\ H \\ \\ H \\ \end{array}$$

$$\begin{array}{c} H \\ \\ H \\ \end{array}$$

$$\begin{array}{c} H \\ \\ H \\ \end{array}$$

$$\begin{array}{c} H \\ \\ H \\ \end{array}$$

- 23. Bases are classified on the basis of their strength, acidity and concentration.
- 24. Strong bases: The bases which almost completely ionized in their aqueous solutions are called strong bases. The solution of strong base contains mostly ions, e.g., NaOH, KOH.
- 25. Weak bases: The bases which incompletely or partially ionized in their aqueous solutions are called weak bases. The solution of weak base contains ions as well as molecules, e.g., NH₄OH, Cu(OH)₂.
- 26. Acidity of a base: The number of replaceable Hydroxyl ions present per molecule of a base in its aqueous solution is called acidity of a base. In case of water insoluble bases, the acidity is equal to the number of Hydroxyl ions present per molecule of a base.

27. Monoacidic bases: The bases which contain only one hydroxyl ion in per molecule of a base.

NaOH
$$\longrightarrow$$
 Na⁺ + OH⁻
KOH \longrightarrow K⁺ + OH⁻
Hydroxyl ion

28. Diacidic bases: The bases which contain two hydroxyl ions in per molecule of a base.

$$Ca(OH)_2 \longrightarrow Ca^{2+} + 2OH^-$$

 $Mg(OH)_2 \longrightarrow Mg^{2+} + 2OH^-$

29. Triacidic bases: The bases which contain three hydroxyl ions in per molecule of a base.

$$Fe(OH)_3 \longrightarrow Fe^{3+} + 3OH^-$$

Ferric hydroxide
$$Al(OH)_3 \longrightarrow Al^{3+} + 3OH^-$$

Aluminium hydroxide

- 30. Methods of preparation of bases: Bases are generally prepared by the following methods:
 - (a) Reaction of metals with oxygen: Metals like sodium, potassium and calcium burn in oxygen to form their respective oxides.

$$4K + O_2 \longrightarrow 2K_2O$$

$$2Ca + O_2 \longrightarrow 2CaO$$

(b) By the reaction of water with soluble metallic oxides and ammonia: Soluble basic oxides and ammonia on dissolving in water produces soluble bases.

$$Na_2O + H_2O \longrightarrow 2NaOH$$

 $K_2O + H_2O \longrightarrow 2KOH$
 $NH_3 + H_2O \longrightarrow NH_4OH$

(c) By the reaction of water on metals: Active metals react with water to displace hydrogen and form metallic hydroxide or metallic oxide depending on the nature of metal and the conditions of temperature.

$$2Na + 2H_2O \longrightarrow 2NaOH + H_2^{\uparrow}$$

$$2K + 2H_2O \longrightarrow 2KOH + H_2^{\uparrow}$$

$$(cold)$$

$$Ca + 2H_2O \longrightarrow Ca(OH)_2 + H_2^{\uparrow}$$

$$(cold)$$

$$Zn + H_2O \longrightarrow ZnO + H_2^{\uparrow}$$

$$(steam)$$

$$3Fe + 4H_2O \Longrightarrow Fe_3O_4 + 4H_2^{\uparrow}$$

$$heated (steam)$$

$$Triferric tetraoxide$$

(d) By the precipitation of metallic hydroxides from their soluble salts: In this case, interchange of radicals in solution takes place, i.e., double decomposition reaction takes place.

(e) Thermal decomposition of metallic carbonates: Metallic carbonates on heating give their respective metallic oxides and carbon-dioxide gas is evolved.

Carbon dioxide is a colourless and odourless gas which turns lime water milky.

$$Ca(OH)_2 + CO_2 \longrightarrow CaCO_3 + H_2O$$

Sodium carbonate and potassium carbonate are stable towards heat.

(f) Thermal decomposition of metallic nitrates: Lead nitrate on heating decripitates, melts to give off reddish brown coloured gas having pungent suffocating smell. A residue is left behind which sticks to the glass tube. The residue is reddish brown or reddish yellow when hot, and light yellow when cold.

$$2Pb(NO_3)_2 \xrightarrow{\Delta} 2PbO + 4NO_2 + O_2$$

$$(Reddish yellow-hot Light yellow-cold)$$
Similarly,
$$2Zn(NO_3)_2 \xrightarrow{\Delta} 2ZnO + 4NO_2 + O_2$$

$$(Yellow - hot White - cold)$$

$$2Cu(NO_3)_2 \xrightarrow{\Delta} 2CuO + 4NO_2 + O_2$$

$$(Black - hot Black - cold)$$

The colour of the residues are summarized in the table given below:

	Color	ur of the residue	
	Name of the residue	Hot	Cold
1.	Lead monoxide	Reddish brown/ Reddish yellow	Light yellow
2.	Zinc oxide	Yellow	White
3.	Copper oxide	Black	Black

Nitrogen dioxide turns freshly prepared acidified ferrous sulphate solution brown black.

$$FeSO_4 + NO_2 \longrightarrow FeSO_4.NO$$

Nitroso ferrous sulphate (Brown black)

Sodium nitrate and potassium nitrate on heating forms their respective nitrite with the liberation of oxygen.

$$2NaNO_{3} \xrightarrow{\Delta} 2NaNO_{2} + O_{2}$$

$$2KNO_{3} \xrightarrow{\Delta} 2KNO_{2} + O_{2}$$

Silver nitrate and mercury nitrate on heating give their respective metal, nitrogen dioxide and oxygen.

$$2AgNO_{3} \xrightarrow{\Delta} 2Ag + 2NO_{2} + O_{2}$$

$$Hg(NO_{3})_{2} \xrightarrow{\Delta} Hg + 2NO_{2} + O_{2}$$

31. The physical properties of bases are

(i) bitter to taste

Ammonium chloride

(ii) slippery to touch.

(iii) Soluble bases show characteristic colour change with indicators.

Name of Indicator	Colour change
1. Red litmus	Blue
2. Phenolphthalein	Pink
3. Methyl orange	Yellow

32. The chemical properties of bases are:

(a) Alkalies react with ammonium salts to liberate a colourless gas having pungent, irritating smell which turns Nessler's reagent brown and give dense white fumes with concentrated HCl.

Ammonium chloride

$$NH_4Cl$$
 + $NaOH \xrightarrow{\Delta} NaCl$ + $NH_3 + H_2O$

Ammonium Sodium Ammonia chloride hydroxide chloride

(b) Alkalies readily absorb carbon dioxide from atmosphere.


(c) Bases react with acids to form salt and water.

(d) Alkalies, i.e., soluble bases precipitate different coloured insoluble bases from their soluble salts.

(e) Insoluble bases decompose on heating to form their respective metallic oxides and water.

$$Zn(OH)_2$$
 $\xrightarrow{\Delta}$ $ZnO + H_2O$
 $Zinc \ hydroxide$ $\xrightarrow{\Delta}$ $CuO + H_2O$
 $Copper \ hydroxide$ $\xrightarrow{\Delta}$ $CuO + H_2O$
 $Copper \ hydroxide$ $\xrightarrow{\Delta}$ $Copper \ oxide$
 $Pb(OH)_2$ $\xrightarrow{\Delta}$ $PbO + H_2O$
 $Lead \ hydroxide$ $Lead \ oxide$


33. pH scale tells us whether the given solution is acidic, alkaline or neutral.



34. pH of pure water is equal to 7.

35. At pH = 7solution is neutral. solution is alkaline. pH > 7, pH < 7, solution is acidic.

36. Universal indicator is a mixture of several organic dyes. It not only tells whether the given solution is acidic or alkaline but it also predicts the strength of the given solution.



Colour changes in the universal indicator (For different pH values)

37. Salts: These are the chemical compounds which on dissolving in water produce positively charged particles other than Hydrogen ions and negatively charged particles other than Hydroxyl ions.

NaCl
$$\rightleftharpoons$$
 Na⁺ + Cl⁻
KCl \rightleftharpoons K⁺ + Cl⁻

- 38. Salts can be classified as
 - (a) Normal salt
- (b) Acid salt (c) Basic salt
- (d) Double salt

- (e) Mixed salt
- (f) Complex salt
- (a) Normal salt: The salt formed by complete replacement of replaceable Hydrogen ions of an acid by a basic radical, or metallic ion. Some examples of normal salts are Sodium chloride (NaCl), Potassium nitrate (KNO3), Copper sulphate (CuSO4), Sodium acetate (CH3COONa), Ammonium chloride (NH₄Cl), Sodium phosphate (Na₃PO₄).

The formation of the above named salts are shown below:

These salts are dissolving in water dissociate into their constituent ions.

$$NaCl \rightleftharpoons Na^+ + Cl^-$$

 $KCl \rightleftharpoons K^+ + Cl^-$

(b) Acid salt: A salt formed by incomplete or partial replacement of replaceable Hydrogen ions of an acid by a basic radical or metallic ion. Some examples of acid salt are Sodium hydrogen sulphate (NaHSO₄), Potassium bisulphate (KHSO₄), Sodium dihydrogen phosphate (NaH₂PO₄), Sodium hydrogen sulphite (NaHSO3). The formation of the above named salts can be shown below.

Acid	+	Alkali	O-++>+*	Salt	+)	Water
H ₂ SO ₄	+	NaOH		NaHSO ₄	+	H_2O
H ₂ SO ₄	4101	КОН	to bus gol a	KHSO ₄	70	H ₂ O
H ₃ PO ₄	+	NaOH		NaH ₂ PO ₄	+	H ₂ O
H ₂ SO ₃		NaOH		NaHSO ₃	+	H_2O

Acid salts are formed when the basicity of the acid taken is more than the acidity of a base. In aqueous solutions, acid salts furnish hydrogen ion or hydronium ion (H3O+). Hence, the aqueous solution of acid salt exhibits all the properties of an acid. The acid salt contains a cation obtained from the base and one or more hydrogn atoms of the acid attached to the anion obtained from the acid.

(c) Basic salt: The salt formed by the partial or incomplete replacement of replaceable Hydroxyl ions of diacidic or triacidic base by an acid radical. Some examples of basic salts are basic lead chloride (Pb(OH)Cl), basic copper chloride (Cu(OH)Cl), Basic magnesium chloride (Mg(OH)Cl), basic lead nitrate (Pb(OH)NO3), basic copper nitrate (Cu(OH)NO). The formation of the above named salts are shown below:

It is to be noted the basic salts are only formed when the acidity of the base taken is more than the basicity of an acid. The basic salts contain a metallic cation, a hydroxyl ion from the base and an anion obtained from the acid.

(d) Double salts: Double salts are formed from two simple salts when they crystallize out from their saturated solutions.

e.g.,
$$K_2SO_4.Al_2(SO_4)_3.24H_2O$$
 - Potash alum
 $FeSO_4.(NH_4)_2.SO_4.6H_2O$ - Mohr's salt
 $K_2SO_4.Cr_2(SO_4)_3.24H_2O$ - Chrome alum
 $(NH_4)_2.SO_4.Fe_2(SO_4)_3.24H_2O$ - Ferric alum

The aqueous solution of double salts furnish the ions of their constituent ions.

(e) Mixed salts: They contain more than one acid or basic radicals.

e.g., $CaOCl_2$ - Bleaching powder $CaOCl_2 \Longrightarrow Ca^{2+} + Cl^- + ClO^-$

CaOCl₂ = Ca²⁺ + Cl⁻ + ClO⁻

NaKCO₃ - Sodium potassium carbonate

(f) Complex salt: They form one complex ion and one simple ion on dissolving in water.

e.g.,
$$K_3[Fe(CN)_6] \rightleftharpoons 3K^+ + [Fe(CN)_6]^{3-}$$

Potassium ion Ferricyanide ion

$$Na[Ag(CN)_2] \rightleftharpoons Na^+ + (Ag(CN)_2]^-$$

Sodium ion Silver cyanide ion

39. Soluble salts are generally prepared by the following methods:

(a) By synthesis: Several soluble salts are prepared by heating the constituent elements together.

(b) Simple displacement: Soluble salts of active metals like Mg, Zn, Fe etc. can be prepared by the simple displacement reactions involving active metal and dilute acid.

(c) Double decomposition - neutralization reaction:

Insoluble or soluble bases react with acids to form salt and water. By this reaction only soluble salts are prepared.

(d) By the reaction of metallic carbonates, bicarbonates, sulphites, bisulphites, sulphides and bisulphides with dilute acids:

KHSO ₃ + HNO ₃ (dil.)	\longrightarrow KNO ₃ + H ₂ O + SO ₂
Potassium	Potassium
bisulphite	nitrate
KHS + HCl (dil.)	\longrightarrow KCl + H_2S
Potassium	Potassium
bisulphide	chloride

(e) By the reaction of chlorides and nitrates with concentrated sulphuric acid:

(f) By the reaction of metals with alkali:

40. Insoluble salts are prepared by the following methods:

(a) Synthesis: Several insoluble salts are prepared by the constituent elements together.

$$Fe + S \xrightarrow{\Delta} FeS$$

$$Iron (II) sulphide$$

$$Zn + S \xrightarrow{\Delta} ZnS$$

$$Zinc sulphide$$

$$Pb + S \xrightarrow{\Delta} PbS$$

$$Lead sulphide$$

(b) Double decomposition - precipitation reaction.

(c) Insoluble salts cannot be directly prepared from another insoluble salt.

These salts are prepared by two steps.

For example conversion of lead carbonate to lead sulphate:

Step I: Conversion of insoluble salt to a soluble salt: By the reaction of insoluble salt with dilute nitric acid.

$$PbCO_3 + 2HNO_3 (dil.) \longrightarrow Pb(NO_3)_2 + H_2O + CO_2$$

Insoluble salt Soluble salt

Step-II: Soluble salt undergoes reaction with either dilute acid or another soluble salt (precipitation).

$$Pb(NO_3)_2 + H_2SO_4 \text{ (dil.)} \longrightarrow PbSO_4 \downarrow + 2HNO_3$$

Soluble salt


$$\begin{array}{c} or \\ \text{Pb(NO}_3)_2 + \text{Na}_2 \text{SO}_4 \xrightarrow{\hspace*{1cm}} \text{PbSO}_4 \downarrow + 2 \text{NaNO}_3 \\ or \end{array}$$

$$Pb(NO_3)_2 + K_2SO_4 \longrightarrow PbSO_4 \downarrow + 2KNO_3$$

41. The laboratory preparation of some normal and acid salts are briefly given below:

(i) Iron (III) chloride or anhydrous ferric chloride (By synthesis)

During the laboratory preparation of anhydrous ferric chloride dry chlorine is passed over heated iron. The iron (III) chloride formed volatizes as its formation is exothermic and it condenses in the receiver as brown scales. As iron (III) chloride is highly deliquescent in nature, therefore it is kept dry in the receiver with the help of anhydrous calcium chloride which does not allow the moisture to enter into the receiver.



Experimental set-up for laboratory preparation of Iron (III) chloride anhydrous

(ii) Copper (II) sulphate (By decomposition)

Copper sulphate can be prepared by the reacion of copper oxide, copper hydroxide or copper carbonate with dilute sulphuric acid.

Add copper oxide or copper hydroxide or copper carbonate to heated dilute sulphuric acid in small quantities with constant strining till no more compound gets dissolved in the acid and the unreacted compound gets settled to the bottom. Filter the hot solution and the filtrate is collected in the China dish. Evaporate the filterate slowly by heating to the point of crystallization and then allow it to cool. Bright blue crystals of CuSO₄.5H₂O are formed which are further dried.

The other salts prepared by this method are lead nitrate, copper nitrate, calcium chloride etc.

(iii) Zinc sulphate (By displacement)

Salts like zinc sulphate, iron (II) sulphate can be prepared by the reaction of metals with dilute sulphuric acid. Metals like iron and zinc can displace hydrogen from dilute acids as they are lying above hydrogen in the metal activity series.

$$\operatorname{Zn} + \operatorname{H_2SO_4}(\operatorname{dil.}) \longrightarrow \operatorname{ZnSO_4} + \operatorname{H_2O}$$
 $\operatorname{Fe} + \operatorname{H_2SO_4}(\operatorname{dil.}) \longrightarrow \operatorname{FeSO_4} + \operatorname{H_2}$

The following steps are involved in the preparation of above named salt

- (i) Dilute sulphuric acid is taken in a beaker and zinc pieces in little quantity are added to it with constant stirring till no further reaction takes place.
- (ii) Effervescence is observed during the liberation of hydrogen. When effervescence stops it indicates the completion of reaction, i.e., all the acid has been used up.
- The above hot solution is filtered and the excess of zinc (unreacted) is removed.
- The filtrate is collected in the China dish and allowed to evaporate to get the crystals.
- (v) The crystals obtained are filtered, washed and dried between the folds of filter paper.

(iv) Sodium sulphate (By neutralization)

Sodium sulphate can be prepared by the reaction of sodium hydroxide or sodium carbonate with dilute sulphuric acid.

$$2\text{NaOH} + \text{H}_2\text{SO}_4 \text{ (dil.)} \longrightarrow \text{Na}_2\text{SO}_4 + 2\text{H}_2\text{O}$$

$$\text{Na}_2\text{CO}_3 + \text{H}_2\text{SO}_4 \text{ (dil.)} \longrightarrow \text{Na}_2\text{SO}_4 + \text{H}_2\text{O} + \text{CO}_2$$

For the preparation of sodium sulphate initially the process of titration is conducted so as to know the volumes of alkali and acid used up for the completion of reaction. As both the reactants and the products are soluble in water. The following steps are involved in the process of titration.

- (i) 5 cm³ of sodium hydroxide solution is measured and transferred in the conical flask. A drop of phenolphhalein is added. The solution turns pink. If Na2CO3 solution is taken then methyl orange is added.
- (ii) Now, rinse and fill the burette with dilute sulphuric acid. Note the initial reading of the burette. Now add acid drop by drop into the conical flask containing sodium hydroxide by constant stirring till the last drop decolourizes the solution (end point-completion of reaction). Now note the final reading of the burette and calculate the difference. The difference will give the volume of dilute sulphuric acid required to neutralize 25 cm3 of sodium hydroxide.
- (iii) The acid and alkali are mixed in the same ratio by volume (as determined by titration) in the evaporating dish. The solution contains sodium sulphate and water.
- (iv) On evaporating the solution the crystals of Na2SO4.10H2O are obtained.

42. The properties of salts are:

(a) Efflorescence: Certain hydrated salts, when left exposed to atmosphere, lose their water of crystallization and crumble into powder. Such salts are called efflorescent salts and the property as efflorescence.

e.g., - $Na_2CO_3.10H_2O$, $Na_2SO_4.10H_2O$, $MgSO_4.7H_2O$

(b) Deliquescence: Certain salts, when left exposed to atmosphere, absorb moisture and get converted into their saturated solution. Such salts are called deliquescent salts and the property as deliquescence.

e.g., - Ferric chloride, Magnesium chloride

(c) Hygroscopy: Certain substances, when left exposed to atmosphere, absorb moisture and become wet. Such substances are called hygroscopic substances and the property as hygroscopy. Hygroscopic substances are called as drying agents.

e.g., - Anhydrous calcium chloride, Calcium oxide, Concentrated sulphuric acid.

(d) Hydrolysis: The phenomenon due to which salt of strong acid and weak base, or weak acid and strong base on dissolving in water reacts with it to form parent acid and alkali thereby rendering the solution either acidic or alkaline in nature.

(i) Ammonium chloride - It is a salt of strong acid and weak base.

Strong acid	Weak base	
Hydrochloric acid (HCl)	Ammonium hydroxide (NH ₄ OH)	
NH ₄ Cl + H ₂ O	NH ₄ OH + HCl	
Highly ionize a NH ⁺ Cl ⁻ + H ₀ O ————————————————————————————————————	Feebly ionized Highly ionized NH ₄ OH + H ⁺ Cl ⁻	

Deleting the spectator ions

$$NH_4^+ + H_2O \longrightarrow NH_4OH + H^+$$

The presence of Hydrogen ions renders the solution acidic.

In general, the salt solution of strong acid and weak base is acidic in nature.

Other examples are Ferric chloride, Copper sulphate, Ammonium sulphate, Ammonium nitrate, Aluminium chloride.

ii) Sodium acetate - CH₃COONa is a salt of strong base, i.e., NaOH and weak acid, i.e., CH₃COOH.

Strong base	Weak acid
Sodium hydroxide (NaOH)	Acetic acid (CH ₃ COOH)

$$CH_3COONa + H_2O \longrightarrow CH_3COOH + NaOH$$
Highly ionized

Weakly ionized

Highly ionized

$$CH_3COO^-Na^+ + H_2O \longrightarrow CH_3COOH + Na^+OH^-$$

Deleting the spectator ions,

$$CH_3COO^- + H_2O \longrightarrow CH_3COOH + OH^-$$

The presence of Hydroxyl ion renders the solution alkaline.

In general, the salt solution of weak acid and strong base is alkaline in nature. Other example, potassium acetate, sodium sulphide, sodium sulphite.

- 43. The salts of strong acid and strong base do not hydrolyse as the solution formed is neutral.
- 44. Water of crystallization: The definite number of water molecules which enters into loose chemical combination when the salt crystallizes out of its saturated solution.
- **45. Hydrated salts on heating** lose their water of crystallization *i.e.*, lose their **crystalline** shape and the **colour fades**., *e.g.*, if hydrated copper sulphate is heated, its colour changes from blue to white and crystalline shape becomes amorphous.

$$CuSO_4$$
. $5H_2O \xrightarrow{\Delta} CuSO_4 + 5H_2O$
Blue, crystalline White, amorphous

- 46. Hydrated salts on heating become anhydrous.
- 47. Zinc and Aluminium displace hydrogen from the acids as well as alkalies, therefore, they are amphoteric in nature.

$$Zn + H_2SO_4 (dil.) \longrightarrow ZnSO_4 + H_2 \uparrow$$

$$Zn + 2HCl (dil.) \longrightarrow ZnCl_2 + H_2 \uparrow$$

$$Zn + 2NaOH \xrightarrow{heat} Na_2ZnO_2 + H_2 \uparrow$$

$$Powder \qquad Sodium$$

$$zincate$$

$$2Al + 3H_2SO_4 (dil.) \longrightarrow Al_2(SO_4)_3 + 3H_2 \uparrow$$

$$2Al + 6HCl(dil.) \longrightarrow 2AlCl_3 + 3H_2 \uparrow$$

$$2Al + 2NaOH + 2H_2O \xrightarrow{heat} 2NaAlO_2 + 3H_2 \uparrow$$

$$Powder \qquad Sodium$$

$$aluminate$$

48. The oxides of Zinc and Aluminium react with acids as well as alkalies to form salt and water as the only products, therefore, they are amphoteric in nature.

$$ZnO + H_2SO_4(dil.) \longrightarrow ZnSO_4 + H_2O$$

$$ZnO + 2HCl(dil.) \longrightarrow ZnCl_2 + H_2O$$

$$ZnO + 2NaOH(dil.) \stackrel{\Delta}{\longrightarrow} Na_2ZnO_2 + H_2O$$

$$Al_2O_3 + 3H_2SO_4(dil.) \longrightarrow Al_2(SO_4)_3 + 3H_2O$$

$$Al_2O_3 + 6HCl(dil.) \longrightarrow 2AlCl_3 + 3H_2O$$

$$Al_2O_3 + 2NaOH(dil.) \stackrel{\Delta}{\longrightarrow} 2NaAlO_2 + H_2O$$

49. The hydroxides of Zinc and Aluminium react with acids as well as alkalies to form salt and water as the only products, therefore, they are amphoteric in nature.

$$\begin{split} &\operatorname{Zn}(\operatorname{OH})_2 + \operatorname{H}_2\operatorname{SO}_4(\operatorname{dil}.) \longrightarrow \operatorname{Zn}\operatorname{SO}_4 + 2\operatorname{H}_2\operatorname{O} \\ &\operatorname{Zn}(\operatorname{OH})_2 + 2\operatorname{Na}\operatorname{OH}\left(\operatorname{dil}.\right) \stackrel{\Delta}{\longrightarrow} \operatorname{Na}_2\operatorname{Zn}\operatorname{O}_2 + 2\operatorname{H}_2\operatorname{O} \\ &2\operatorname{Al}(\operatorname{OH})_3 + 3\operatorname{H}_2\operatorname{SO}_4(\operatorname{dil}.) \longrightarrow \operatorname{Al}_2(\operatorname{SO}_4)_3 + 6\operatorname{H}_2\operatorname{O} \\ &\operatorname{Al}(\operatorname{OH})_3 + \operatorname{Na}\operatorname{OH}(\operatorname{dil}.) \stackrel{\Delta}{\longrightarrow} \operatorname{Na}\operatorname{Al}\operatorname{O}_2 + 2\operatorname{H}_2\operatorname{O} \end{split}$$

50. Solubility Table of Salts

- (i) All salts of Sodium, Potassium and Ammonium are soluble in water.
- (ii) All nitrates and nitrites are soluble in water.
- (iii) All sulphates are soluble in water. Except: CaSO, PbSO, BaSO,
- (iv) All chlorides are soluble in water.
 - Except: Hg2Cl2, AgCl and PbCl2 (soluble in hot water)
- (v) All sulphites are insoluble in water. Except: Na,SO, K,SO, and (NH,)2SO,
- (vi) All sulphides are insoluble in water Except: Na,S, K,S and (NH,),S.
- (vii) All carbonates are insoluble in water. Except: Na, CO, K, CO, and (NH,), CO,

B-IZaCO - Illibligoide (III) Mer) - (AN O-

51. Equation for the acid rain: The oxides of sulphur (released from oil refineries) and the oxides of nitrogen (released by automobiles exhaust) produces acid rain.

Sulphurdioxide combines with oxygen and water to form sulphuric acid.

$$2SO_2 + 2H_2O + O_2 \longrightarrow 2H_2SO_4$$

Nitrogen dioxide form nitric acid.

$$N_2 + O_2 \xrightarrow{Thunder and} 2NO$$

$$2NO + O_2 \xrightarrow{Dightning} 2NO_2$$

$$4NO_2 + 2H_2O + O_2 \xrightarrow{Dightning} 4HNO_3$$

dilute bydrochieric and

(C) Sulphurtrioxele

one substance in each case which matches the description given in parts (i) to (iv). (Note: Each substance is used only once (i) Sodium thiosulphate is reacted with in the answer.)

(A) Iron(III) chloride

(B) Chromium sulphate

(C) Lead(II) chloride

(D) Sodium chloride.

dilute hydrochloric acid.

hydrochloric acid.

sodium sulphite.

(ii) Calcium bicarbonate reacts with dilute

(iii) Dilute sulphuric acid is poured over

- (i) A compound which is deliquescent.
- (ii) A compound which is insoluble in cold water, but soluble in hot water.
- (iii) A compound whose aqueous solution is neutral in nature.
- (iv) The compound which is responsible for the green colouration when sulphur dioxide is passed through acidified potassium dichromate solution. [5]

(iii) D (ii)

Q2. Select the correct answer from the choices A, B, C and D which are given. Write only the letter corresponding to the correct answer.

- (i) An organic weak acid is:
 - (A) Formic acid
 - (B) Sulphuric acid
 - (C) Nitric acid
 - (D) Hydrochloric acid

Ans. A

- (ii) An example of a complex salt is:
 - (A) Zinc sulphate
 - (B) Sodium hydrogen sulphate
 - (C) Iron(II) ammonium sulphate
 - (D) Tetraammine copper(II) sulphate. [1]

Ans. D

Q3. State your observation for the following cases:


Paper soaked in potassium permanganate solution is introduced into a gas jar of sulphur dioxide.

Ans. It decolourizes purple colour of KMnO₄.

Q4. Write the equation for the reactions: Zinc oxide is treated with sodium hydroxide solution.

Ans. $ZnO + 2NaOH \longrightarrow Na_2ZnO_2 + H_2O$

Q5. Give the equations for the following conversions A to E.



Ans. A:
$$ZnSO_4 + Na_2CO_3 \longrightarrow ZnCO_3 + Na_2SO_4$$
.
B: $ZnCO_3 + 2HNO_3 \longrightarrow Zn(NO_3)_2 + H_2O + CO_2$.

- $C: Zn(NO_3)_2 + 2NaOH \longrightarrow Zn(OH)_2 +$ 2NaNO₃.
- $D: Zn(OH)_2 \xrightarrow{\Delta} ZnO + H_2O.$
- $E: ZnO + H_2SO_4 \longrightarrow ZnSO_4 + H_2O$
- Q6. Solution A is a sodium hydroxide solution. Solution B is a weak acid. Solution C is dilute sulphuric acid. Which solution will
 - (i) liberate sulphur dioxide from sodium sulphite.
 - (ii) give a white precipitate with zinc sulphate solution.
 - (iii) contain solute molecules and ions? [3]

Ans. (i) C (ii) A (iii) B

- Q7. Give the equation for the preparation of each of the following salts from the starting material given.
 - (i) Copper sulphate from copper(II) oxide.
 - (ii) Iron(III) chloride from Iron.
 - (iii) Potassium sulphate from potassium hydroxide solution.
 - (iv) Lead chloride from lead carbonate [5] (two equations).
- Ans. (i) $CuO + H_2SO_4 \longrightarrow CuSO_4 + H_2O$.


(ii) $2\text{Fe} + 3\text{Cl}_2 \longrightarrow 2\text{FeCl}_3$.

(iii) 2KOH + $H_2SO_4 \longrightarrow K_2SO_4 + 2H_2O$

 $(iv) \text{ PbCO}_3 + 2\text{HNO}_3 \longrightarrow \text{Pb(NO}_3)_2 +$ $H_9O + CO_9$ $Pb(NO_3)_2 + 2HCl \longrightarrow PbCl_2 + 2HNO_3$.

Q1. (a) The diagram given below is to prepare Iron(III) chloride in the laboratory:

2009



- What is substance B?
- (ii) What is the purpose of B?
- (iii) Why is iron(III) chloride to be stored in a closed container?

	(iv) Write th	e equation	for the
	reaction	between i	iron and
	chlorine.		[4]
(b) W	rite the ed	quation(s)	for the
and plot re	action(s) to p	repare lead	sulphate
fr	om lead carbo	nate.	[2]
Ans. (a) (i	Anhydrous ca	lcium chlorid	e
	Drying agent		(1)
	As it is highly		11163
	$2Fe + 3Cl_2$	THE RESERVE OF THE PARTY OF THE	(1)
	O ₃ + 2HNO ₃ (dil	THE RESERVE OF THE PARTY OF THE	O) + H O
(0)150	03 1 2111103 (dil	71 0(11	The second second
	tion and the same of		+ CO ₂
THE RESERVE THE PROPERTY OF THE PERSON NAMED IN COLUMN TWO IN COLUMN TO THE PERSON NAMED IN COLU	$(0_3)_2 + H_2SO_4$ (dil.		*
	the term: Ne		
	H ⁺ ions derived		
	ydroxyl ions of a		
water	molecule is calle	ed as neutran	sation.
	2008	- AUGUSTO	NO.
Q1. Select	the correct an	swer from th	e choices
	, D which are		ic choices
	sulphuric aci	The second second second second	e a white
	itate when ad		
	pper nitrate		Ams. (5) (
B. Zin	c nitrate		
C. Lea	ad nitrate		
D. Soc	dium nitrate		[1]
Ans. C			
	re the terms de		
	salt contai		
	rounded by ot		
VI 750 CHE	pase which is s		
Ans. (i) Cor			
(ii) Alk			
	g use only of su given below :	ibstances che	osen from
	lphuric acid	Sodium ca	whomato
Zinc	iphuric acid	Sodium su	
Lead		Calcium ca	
	he equations		
	you could obt		
100000 000	drogen		
(ii) sul	phur dioxide		
(iii) car	bon dioxide		
(iv) zin	c carbonate (t	wo steps req	uired) [5]
	+ H ₂ SO ₄ (dil.) —		
(ii) Na ₂	$SO_3 + H_2SO_4(dil.)$	\longrightarrow Na ₂ SO ₄ +1	$H_2O + SO_2\uparrow$

(iii) Na ₂ CO ₃ + H ₂ S		
(iv) Zn + H ₂ SO ₄ (di		2 4
$ZnSO_4 + Na_2C$	$O_3 \longrightarrow ZnC$	$O_3 + Na_2SO_4$
maitata attituta attituta	2007	B ATOMPO
Q1. From the list	given belo	w. select th
word(s) requir		
blanks (i) to (v)		373
ammonia, a	ammonium	, carbonate
carbon dioxi		
hydroxide, p	recipitate, sa	alt, water.
A solution X		
it must conta		
another solu		
blue and th		
(ii) X and Y ar		
products will		
and (iv)		
magnesium v		
(v)	_ gas would	be evolved.
		[5]
(Note: Words		
to be used onl		
as (i), (ii), (iii) and so on.	Do not copy
the passage.)	(ii) Undannid	d sens
Ans. (i) Hydronium (iii) Salt	(iv) Water	
(v) Hydrogen	(iv) water	
Q2. Match the follow	ing	solutios
Column A	and the latest and the same	umn B
1. Acid salt		n potassium
for time reactions:	carbon	A CONTRACTOR OF THE PERSON NAMED IN CONT
2. Mixed salt	B. Alum	
3. Complex salt		n carbonate
4. Double salt	D. Sodiur	
5. Normal salt		n hydrogen
Ans. 1. Acid salt :	carbon	
	A STATE OF THE PARTY OF THE PARTY.	gen carbonate
2. Mixed salt : S 3. Complex salt :	CONTRACTOR OF THE PERSON NAMED IN	
4. Double salt :		Le (I) to (In).
5. Normal salt : S		ato
Q3. Write balance		
following reaction	- cquation	is for the

following reactions:

(i) Lead sulphate from lead nitrate

solution and dilute sulphuric acid.

- (ii) Copper sulphate from copper and concentrated sulphuric acid.
- (iii) Lead chloride from lead nitrate solution and sodium chloride solution.
- (iv) Ammonium sulphate from ammonia and dilute sulphuric acid.
- (v) Sodium chloride from sodium carbonate solution and dilute hydrochloric acid. [5]
- Ans. (i) $Pb(NO_3)_2 + H_2SO_4 \longrightarrow PbSO_4 + 2HNO_3$
 - (ii) Cu + 2H₂SO₄(conc.) \longrightarrow CuSO₄ + 2H₂O + SO₂
 - (iii) Pb(NO₃)₂ + 2NaCl \longrightarrow PbCl₂ + 2NaNO₃
 - (iv) 2NH₃ + H₂SO₄(dil.) \longrightarrow (NH₄)₂SO₄
 - (v) $Na_2CO_3 + 2HCl(dil.) \longrightarrow 2NaCl + H_2O + CO_2$
- Q4. Salts A, B, C, D and E undergo reactions (i) to (v) respectively. Identify the anion present in these salts on the basis of these reactions. Tabulate your answers in the format given below:
 - (i) When silver nitrate solution is added to a solution of A, a white precipitate, insoluble in dilute nitric acid, is formed.
 - (ii) Addition of dilute hydrochloric acid to B produces a gas which turns lead acetate paper black.
 - (iii) When a freshly prepared solution of ferrous sulphate is added to a solution of C and concentrated sulphuric acid is gently poured from the side of the test tube, a brown ring is formed.
 - (iv) When dilute sulphuric acid is added to D a gas is produced which turns acidified potassium dichromate solution from orange to green.
 - (v) Addition of dilute hydrochloric acid to E produces an effervescence. The gas produced turns lime water milky but does not affect acidified potassium dichromate solution.

PAU	Salt	ribor	Anion	Date on	
B.A.	A	37/2	al grange (Split	1057	
Mil Y	В	BALLES.		Charles	
1000	C	STATE OF		Euch :	
105	D	The same		STATE OF	
	E		STATE OF THE PARTY		[5

- Ans. (i) A Chloride or Cl
 - (ii) B-Sulphide or S2-
 - (iii) C Nitrate or NO₃
 - (iv) D Sulphite or SO_3^{2-}
 - (v) E Carbonate or CO_3^{2-}

-2006

- Q1. Mention the colour changes observed when the following indicators are added to acids:
 - (i) Alkaline phenolphthalein solution
 - (ii) Methyl orange solution
 - (iii) Neutral litmus solution [3]

Ans. (i) It gets decolourised

- (ii) It turns red or pink
- (iii) It turns red.
- Q2. Choose the correct answers from the choices A, B, C and D.

Which of the following hydroxides is not an alkali.

- A: Ammonium hydroxide
- B: Calcium hydroxide
- C: Copper hydroxide
- D: Sodium hydroxide [3]

Ans. C: Copper hydroxide

2005_

- Q1. Write balanced chemical equations for the following:
 - (i) Potassium hydrogen carbonate and dilute sulphuric acid
- (ii) Copper oxide and dilute hydrochloric acid [2]
- Ans. (i) KHCO₃ + H₂SO₄(dil.) \longrightarrow KHSO₄ + H₂O + CO₂
 Or

 $2KHCO_3 + H_2SO_4(dil.) \rightarrow K_2SO_4 + 2H_2O + 2CO_2$

- (ii) $CuO + 2HCl(dil.) \longrightarrow CuCl_2 + H_2O$
- Q2. The questions (i) to (v) refer to the following salt solutions listed A to F:

A: Copper nitrate

B: Iron (II) sulphate

C: Iron (III) chloride

D: Lead nitrate

E: Magnesium sulphate

F: Zinc chloride

- (i) Which two solutions will give a white precipitate when treated with dilute hydrochloric acid followed by barium chloride solution?
- (ii) Which two solutions will give a white precipitate when treated with dilute nitric acid followed by silver nitrate solution?
- (iii) Which solution will give a white precipitate when either dilute hydrochloric acid or dilute sulphuric acid is added to it?
- (iv) Which solution becomes deep / inky blue colour when excess ammonium hydroxide is added to it?
- (v) Which solution gives a white precipitate with excess ammonium hydroxide solution? [5]
- Ans. (i) Iron (II) sulphate, Magnesium sulphate
 - (ii) Iron (III) chloride, Zinc chloride
 - (iii) Lead nitrate
 - (iv) Copper nitrate
 - (v) Lead nitrate
 - Q3. Match the description given below:
 - A : Acidic oxide F : Efflorescence
 - B : Alkali G : Electrolysis
 - C : Amphoteric H : Electrolyte oxide
 - D : Basic oxide I : Homologous series
 - E: Deliquescence J: Hydrocarbons
 - (i) The property of spontaneously giving up water of crystallization to the atmosphere.
 - (ii) A compound, which is soluble in water and the only negative ions in the solution are hydroxide ions.
 - (iii) An oxide, which forms salts when it reacts with both acids and alkalies. [3]
- Ans. (i) Efflorescence (ii) Alkali
 - (iii) Amphoteric oxide

- Q4. What do you observe when
 - (i) Hydrogen sulphide gas is passed through Lead acetate solution?
 - (ii) Neutral litmus solution is added to Sodium hydrogen carbonate solution?

Ans. (i) A black precipitate appears

(ii) It turns Red

- Q5. Fill in the blanks with suitable words.
 - An acid is a compound which when dissolved in water forms hydronium ions as the only (1) _ _ _ ions. A base is a compound which if soluble in water contains (2) _ _ _ ions. A base reacts with an acid to form a (3) _ _ _ and water only. The type of reaction is known as (4) _ _ _ [4]

Ans. 1. Positive 2. Hydroxyl

3. Salt 4. Neutralisation

- Q6. The preparation of lead sulphate from lead carbonate is a two step process (Lead sulphate cannot be prepared by adding dilute sulphuric acid to lead carbonate).
 - (i) What is the first step that is required to prepare lead sulphate from lead carbonate?
 - (ii) Write the equation for the reaction that will take place when this first step is carried out?
 - (iii) Why is the direct addition of dilute sulphuric acid to lead carbonate an impractical method of preparing lead sulphate? [3]
- Ans. (i) The first step is to convert lead carbonate to lead nitrate by reacting it with dilute Nitric acid.
 - (ii) $PbCO_3 + 2HNO_3 \longrightarrow Pb(NO_3)_2 + H_2O + CO_2$
 - (iii) The insoluble lead sulphate formed does not allow the reaction between acid and the salt.

IMPORTANT QUESTIONS

- Q1. (a) (i) What is the purpose of pH scale?
 - (ii) What is the pH of pure water?
 - (iii) 'A' is a soluble acidic oxide, 'B' is a soluble base. Compared to the pH of pure water, what will be the pH of (1) a solution of 'A' (2) a solution of 'B'
- (b) Taking Sodium carbonate as an example, give the meaning of the following terms:
 - (i) Water of Crystallization
 - (ii) Anhydrous
 - (iii) Efflorescence

- (c) Barium chloride solution can be used to distinguish between a Sodium sulphate solution and a Sodium nitrate solution. How is this done?
- (d) Name two gases which are responsible for acid rain.
- Ans. (a) (i) pH scale is used to determine the alkalinity and acidic nature of the solution.
 - (ii) Seven
 - (iii) (1) Less than seven (2) More than seven
 - (b) (i) Water of Crystallization: The definite number of water molecules which enter into loose chemical combination when he salt crystallizes out from its saturated solution.

Sodium carbonate ($Na_2CO_3.10H_2O$) has ten water molecules of crystallization.

(ii) Anhydrous: Hydrated salts on heating lose their water of crystallization and are rendered anhydrous.

$$Na_2CO_3.10H_2O \xrightarrow{\Delta} Na_2CO_3 + 10H_2O$$
 anhydrous

- (iii) Efflorescence: Hydrated salts, when left exposed to atmosphere, lose their water of Crystallization and crumble down to form a powder. Such salts are called as efflorescent salts and the property as efflorescence.
 - Na₂CO₃.10H₂O, on exposure to atmosphere, loses its water of Crystallization and crumbles to form a powder.
- (c) Barium chloride solution, when reacts with Sodium sulphate solution, a white precipitate is formed which is insoluble in all the mineral acids, whereas when Barium chloride solution reacts with Sodium nitrate, no visible reaction takes place.

 $BaCl_2 + Na_2SO_4 \longrightarrow BaSO_4 + 2NaCl$ $BaCl_2 + 2NaNO_3 \longrightarrow Ba(NO_3)_2 + 2NaCl$

- (d) Nitrogen dioxide and Sulphur dioxide.
- Q2. (a) Outline the steps required to convert Hydrogen chloride to anhydrous Iron (III) chloride. Write the equations for the reactions which take place.
 - (b) (i) What are the two steps to change Lead carbonate to Lead chloride?
 - (ii) Give the name of a soluble salt of lead and write the equation for the action of heat on this salt.

Ans. (a) MnO₂+4HCl(conc.) → MnCl₂+Cl₂+2H₂O
Chlorine is dried by passing through
concentrated H₂SO₄. Then dried Chlorine
is passed over heated Iron to get anhydrous
Iron (III) chloride.

 $2Fe + 3Cl_2 \longrightarrow 2FeCl_3$

- (b) (i) $PbCO_3+2HNO_3(dil.) \rightarrow Pb(NO_3)_2+H_2O+CO_2$ $Pb(NO_3)_2+2HCl(dil.) \rightarrow PbCl_2+2HNO_3$
 - (ii) Lead nitrate

 $2Pb(NO_3)_2 \xrightarrow{\Delta} 2PbO + 4NO_3 + O_2$

Q3. From the following list of substances, choose those which meet the description given below:

Ammonium chloride, Ammonium nitrate, Chlorine, dilute Hydrochloric acid, Iron, Lead nitrate, Manganese (IV) oxide, Silver nitrate, Sodium nitrate, Sulphur.

Two compounds whose aqueous solution gives white precipitate with dilute Hydrochloric acid.

- Ans. Silver nitrate solution and Lead nitrate solution.
- Q4. What do you see when Barium chloride solution is added to dilute Sulphuric acid?
- Ans. A white precipitate is formed which is insoluble in all mineral acids.
 - Q5. (a) A solution has a pH of 7. Explain how you would
 - (i) increase its pH.
 - (ii) decrease its pH.
 - (b) If a solution changes the colour of litmus from red to blue, what can you say about its pH?
 - (c) What can you say about the pH of a solution that liberates Carbon dioxide from Sodium carbonate?
- Ans. (a) (i) By adding alkali
 - (ii) By adding an acid.
 - (b) More than seven.
 - (c) Less than seven.
 - Q6. Write equations for each of the following reactions:
 - (a) Chlorine is passed over heated iron.
 - (b) Copper sulphate solution is added to Sodium hydroxide solution.

Ans. (a) $2\text{Fe} + 3\text{Cl}_2 \longrightarrow 2\text{FeCl}_3$ heated dry

(b) $CuSO_4 + 2NaOH \longrightarrow Cu(OH)_2 + Na_2SO_4$

Q7. Name from the list of substances given below, the substances which you would use to prepare each of the following salts named in parts (a) to (d):

Copper, Lead, Sodium, Zinc, Copper oxide, Lead carbonate, Sodium carbonate solution, dilute Hydrochloric acid, dilute Nitric acid and dilute Sulphuric acid.

- (a) Zinc sulphate
- (b) Copper sulphate
- (c) Sodium sulphate
- (d) Lead sulphate

Ans. (a) $Zn + H_2SO_4(dil.) \longrightarrow ZnSO_4 + H_2 \uparrow$

- (b) $CuO + H_2SO_4(dil.) \longrightarrow CuSO_4 + H_2O$
- (c) $Na_2CO_3 + H_2SO_4(dil.) \longrightarrow Na_2SO_4 + H_2O + CO_2$
- (d) $PbCO_3 + 2HNO_3(dil.) \longrightarrow Pb(NO_3)_2 + H_2O + CO_2$ $Pb(NO_3)_2 + H_2SO_4(dil.) \longrightarrow PbSO_4 + 2HNO_3$
- Q8. For each of the salts A, B, C and D, suggest a suitable method of preparation which relates to its description given below:
 - (a) 'A' is Sodium salt.
 - (b) 'B' is an insoluble salt.
 - (c) 'C' is a soluble salt of Copper.
 - (d) 'D' is a soluble salt of Zinc.
 - (Do not describe the procedure for the preparation. Refer only to the appropriate method)
- Ans. (a) By neutralization
 - e.g., NaOH + HCl \longrightarrow NaCl + H₂O
 - (b) By precipitation
 - $e.g., Pb(NO_3)_2 + 2NaCl \longrightarrow PbCl_2 \downarrow + 2NaNO_3$
 - (c) $CuCO_3 + H_2SO_4(dil.) \longrightarrow CuSO_4 + H_2O + CO_2$
 - (d) Simple displacement
 - e.g., $Zn + H_2SO_4(dil.) \longrightarrow ZnSO_4 + H_2\uparrow$
- Q9. Define or explain the meaning of the following terms:
 - (a) Acid (b) pH scale.
- **Ans.** (a) Acid: Acid is the chemical compound which, on dissolving in water, produces Hydronium ions as the only positively charged particles.
 - (b) pH scale: The scale used for determining the acidic or alkaline character of the solution.


- Q10. Sulphuric acid can be used to prepare a number of gases in the laboratory. Write balanced equations for reactions in which the following gases are obtained using dilute Sulphuric acid as one of the reactants.
 - (a) Hydrogen
 - (b) Carbon dioxide
 - (c) Sulphur dioxide
- Ans. (a) $Zn + H_2SO_4(dil.) \longrightarrow ZnSO_4 + H_2 \uparrow$ or any other active metal except Na, K, Ca, Sn, Pb.
 - (b) Na₂CO₃+H₂SO₄(dil.)→Na₂SO₄+H₂O+ CO₂↑ any other metallic Carbonate or Bicarbonate except of Calcium and Lead.
 - (c) $Na_2SO_3 + H_2SO_4(dil.) \longrightarrow Na_2SO_4 + H_2O + SO_2 \uparrow$ Any other metallic Sulphite or Bisulphite.
- Q11. (a) The pH value of pure water is 7.

 Compare the pH values of Sulphur dioxide solution and Ammonia solution with that of pure water.
 - (b) Name:
 - (i) two bases which are not alkalies.
 - (ii) a normal salt and acid salt of same acid.
 - (iii) a salt insoluble in cold water but soluble in hot water.
- Ans. (a) pH of Sulphur dioxide solution is less than seven.

pH of Ammonia solution is more than seven.

- (b) (i) Magnesium hydroxide, Copper hydroxide.
 - (ii) Sodium sulphate and Sodium bisulphate.
 - (iii) Lead chloride.
- Q12. Answer the following questions relating to salts and their preparation:
 - (a) What is salt?
 - (b) What kind of salt is prepared by precipitation?
 - (c) Name a salt prepared by direct combination. Write the equation for the reaction that takes place when preparing the salt you have named.
 - (d) What procedure would be used to prepare Sodium salt such as Sodium sulphate. (Give the name of the procedure only)

- Ans. (a) Salts are the chemical compounds which, on dissolving in water, produce positively charged particles other than Hydronium ions and negatively charged particles other than Hydroxyl ion.
 - (b) Insoluble salts are prepared by precipitation.
 - (c) Iron (III) chloride or Ferric chloride $2Fe + 3Cl_2 \longrightarrow 2FeCl_3$ heated dry
 - (d) By reacting Sodium carbonate or Sodium hydroxide with dilute Sulphuric acid. It is prepared by the reaction of neutralization.
 Na₂CO₃+H₂SO₄(dil.)—Na₂SO₄+H₂O+CO₂
 2NaOH + H₂SO₄(dil.)—Na₂SO₄ + 2H₂O
- Q13. For each of the conversions A, B, C, D and E in the scheme below, state briefly in words how the conversions can be carried out:




Ans. A. $CuSO_4 + Na_2CO_3 \longrightarrow CuCO_3 + Na_2SO_4$

- B. $CuCO_3 \xrightarrow{\Delta} CuO + CO_2 \uparrow$
- C. $CuSO_4 + 2NaOH \longrightarrow Cu(OH)_2 + Na_2SO_4$
- D. $Cu(OH)_2 \longrightarrow CuO + H_2O$
- E. $CuO + H_2SO_4 \longrightarrow CuSO_4 + H_2O$
- Q14. (i) What is an alkali?
 - (ii) Explain the meaning of the term acid salt.
- Ans. (i) Water soluble bases are called as alkalies.
 - (ii) The salt formed by the incomplete replacement of replaceable Hydrogen ions of an acid by a basic radical or metallic ion.
- Q15. State what would you observe when:
 - (a) Hydrated Copper sulphate is heated.
 - (b) Sodium carbonate solution is mixed with Calcium chloride solution.
 - (c) Lead nitrate solution is mixed with dilute Hydrochloric acid.
 - (d) Lead nitrate is heated.
 - (e) Ferric chloride is exposed to atmosphere.
 - (f) Blue litmus is dipped in the solution of dilute acid.

- (g) Dilute acid is added to Sodium carbonate.
- (h) Sodium hydroxide solution is slowly added to Copper sulphate solution.
- (i) Copper hydroxide is heated.
- (j) Sodium hydroxide is added to Ammonium chloride.
- (k) Copper nitrate is heated.
- (l) Copper carbonate is heated.
- (m) Barium chloride solution is slowly added to Sodium sulphate solution.
- (n) Carbon dioxide is passed through lime water.
- (o) Washing soda crystals are exposed to atmosphere.
- Ans. (a) Hydrated Copper sulphate, on heating, changes from blue to white and the crystalline form changes to amorphous (powder).
 - (b) A white precipitate is observed.
 - (c) A white precipitate is formed which is soluble in hot water but insoluble in cold water.
 - (d) Lead nitrate, on heating, decripitates, melts to give a reddish brown coloured gas which turns freshly prepared acidified Ferrous sulphate solution brown black. A residue is left behind which is reddish brown when hot and light yellow when cold.
 - (e) It absorbs moisture from the atmosphere and gets converted into its solution.
 - (f) It changes to red.
 - (g) A colourless and odourless gas with brisk effervescence is evolved which turns lime water milky.
 - (h) A bluish white precipitate is formed which is insoluble in excess of Sodium hydroxide.
 - (i) Bluish white precipitate slowly changes to black colour.
 - (j) A colourless gas with pungent irritating smell evolves which turns Nessler's reagent brown and gives dense white fumes with concentrated Hydrochloric acid.
 - (k) Copper nitrate, on heating, melts to give a reddish brown coloured gas which turns freshly prepared acidified Ferrous sulphate solution brown black. A residue is left behind which is black when hot and black when cold.

- (l) Copper carbonate, on heating, gives a colourless and odourless gas which turns lime water milky. A residue is left behind which is black when hot and black when cold.
- (m) A white precipitate is formed which is insoluble in all mineral acids.
- (n) It turns milky.
- (o) It crumbles to form a powder.
- Q16. How are the following conversions carried out? Give balanced chemical equations only.



$$(c) \text{ Zn} \xrightarrow{A} \text{ZnS} \xrightarrow{B} \text{ZnSO}_{4} \xrightarrow{C} \text{Zn(OH)}_{2} \xrightarrow{D} \text{ZnO}$$

ZnCO₃

(e) $Fe^{-A} \rightarrow FeCl_3 \xrightarrow{B} Fe(OH)_3 \xrightarrow{C} Fe_2O_3$

H₂S

Ans. (a) A: CuO + $H_2SO_4(dil.)$ \longrightarrow CuSO₄ + H_2O B: CuSO₄+2NaOH \longrightarrow Cu(OH)₂ + Na₂SO₄

 $C: Cu(OH)_2 \xrightarrow{\Delta} CuO + H_2O$

D: $CuO + 2HNO_3(dil.) \longrightarrow Cu(NO_3)_2 + H_2O$

E: $2Cu(NO_3)_2 \xrightarrow{\Delta} 2CuO + 4HNO_2 + O_2 \uparrow$ F: $CuSO_4 + Na_2CO_3 \longrightarrow CuCO_3 + Na_2SO_4$

 $G: CuCO_3 \xrightarrow{\Delta} CuO + CO_9$

(b) A: PbO+2HNO₃(dil.) \longrightarrow Pb(NO₃)₂ + H₂O B: Pb(NO₃)₂+2NaOH \rightarrow Pb(OH)₂ + 2NaNO₃

 $C: Pb(OH)_2 \xrightarrow{\Delta} PbO + H_2O$

D: $2Pb(NO_3)_2 \xrightarrow{\Delta} 2PbO + 4NO_2 + O_2 \uparrow$ E: $Pb(NO_3)_2 + 2NaCl \longrightarrow PbCl_2 + 2NaNO_3$

(c) A : $Zn + S \xrightarrow{\Delta} ZnS$

 $B : ZnS + H_2SO_4(dil.) \longrightarrow ZnSO_4 + H_2S$

C: $ZnSO_4 + 2NaOH \longrightarrow Zn(OH)_2 + Na_2SO_4$

 $D: Zn(OH)_2 \xrightarrow{\Delta} ZnO + H_2O$

 $E : ZnS + 2HCl(dil.) \longrightarrow ZnCl_2 + H_2S$

F : ZnSO₄+Na₂CO₃ --- ZnCO₃+Na₂SO₄

 $G: ZnCO_3 \xrightarrow{\Delta} ZnO + CO_2$

(d) A : Fe + S $\xrightarrow{\Delta}$ FeS

B : FeS + 2HCl(dil.) → FeCl₂ + H₂S

 $C : FeCl_2 + 2NaOH \longrightarrow Fe(OH)_2 + 2NaCl$

D: $Fe(OH)_2 \xrightarrow{\Delta} FeO + H_2O$

 $E : FeS + 2HCl \longrightarrow FeCl_2 + H_2S$

 $F : FeCl_2 + Na_2CO_3 \longrightarrow FeCO_3 + 2NaCl$

 $G : FeCO_3 \xrightarrow{\Delta} FeO + CO_2$

(e) A : 2Fe + 3Cl₂ → 2FeCl₃
heated dry

B: $FeCl_3 + 3NaOH \longrightarrow Fe(OH)_3 + 3NaCl$

C: $2\text{Fe}(OH)_3 \xrightarrow{\Delta} \text{Fe}_2O_3 + 3H_2O$

Q17. How can you obtain the following gases by using dilute acid and one another substance?

(a) Hydrogen

(b) Carbon dioxide

(c) Sulphur dioxide

(d) Hydrogen sulphide

Ans. (a) Fe + $H_2SO_4(dil.) \longrightarrow FeSO_4 + H_2\uparrow$ $Mg + 2HCl(dil.) \longrightarrow MgCl_2 + H_2\uparrow$

> (b) $Na_2CO_3+2HCl(dil.) \longrightarrow 2NaCl + H_2O + CO_2 \uparrow$ $NaHCO_3+HCl(dil.) \longrightarrow NaCl + H_2O + CO_2 \uparrow$

> (c) $Na_2SO_3+2HCl(dil.)\longrightarrow 2NaCl+H_2O+SO_2\uparrow$ $Na_2SO_3+2H_2SO_4(dil.)\longrightarrow Na_2SO_4+H_2O+SO_2\uparrow$

(d) FeS + 2HCl(dil.) \longrightarrow FeCl₂ + H₂S \uparrow Na₂S + 2HCl(dil.) \longrightarrow 2NaCl + H₂S \uparrow ZnS + H₂SO₄(dil.) \longrightarrow ZnSO₄ + H₂S \uparrow

Q18. How are the following named salts prepared in laboratory? Give balanced chemical equations only.

(a) Iron (III) chloride

(b) Magnesium sulphate

(c) Sodium sulphate

(d) Lead chloride

- (e) Lead nitrate
- (f) Iron (II) sulphide
- (g) Sodium chloride
- (h) Iron (II) chloride
- (i) Zinc sulphate
- (j) Lead sulphide
- (k) Lead carbonate
- (l) Lead sulphate
- (m) Copper sulphate
- (n) Magnesium chloride
- (o) Silver chloride
- Ans. (a) $2\text{Fe} + 3\text{Cl}_2 \longrightarrow 2\text{FeCl}_3$ heated dry
 - (b) $Mg + H_2SO_4(dil.) \longrightarrow MgSO_4 + H_2 \uparrow$
 - (c) $Na_2CO_3+H_2SO_4(dil.)$ $\longrightarrow Na_2SO_4+H_2O+CO_2$ \uparrow $2NaOH+H_2SO_4(dil.)$ $\longrightarrow Na_2SO_4+2H_2O$
 - (d) $Pb(NO_3)_2 + 2HCl(dil.) \longrightarrow PbCl_2 \downarrow + 2HNO_3$ $Pb(NO_3)_2 + 2NaCl \longrightarrow PbCl_2 \downarrow + 2NaNO_3$
 - (e) $PbO + 2HNO_3(dil.) \longrightarrow Pb(NO_3)_2 + H_2O$
 - (f) $Fe + S \xrightarrow{\Delta} FeS$
 - (g) $2\text{Na} + \text{Cl}_2 \longrightarrow 2\text{NaCl}$ $\text{NaOH} + \text{HCl(dil.)} \longrightarrow \text{NaCl} + \text{H}_2\text{O}$
 - (h) Fe + 2HCl(dil.) \longrightarrow FeCl₂ + H₂ \uparrow
 - (i) $\operatorname{Zn} + \operatorname{H}_2\operatorname{SO}_4(\operatorname{dil}.) \longrightarrow \operatorname{ZnSO}_4 + \operatorname{H}_2 \uparrow$ $\operatorname{ZnO} + \operatorname{H}_2\operatorname{SO}_4(\operatorname{dil}.) \longrightarrow \operatorname{ZnSO}_4 + \operatorname{H}_2\operatorname{O}$
 - (j) $Pb(NO_3)_2 + H_2S \longrightarrow PbS + 2HNO_3$ $Pb(NO_3)_2 + Na_2S \longrightarrow PbS + 2NaNO_3$
 - $(k) \text{ Pb(NO}_3)_2 + \text{Na}_2\text{CO}_3 \longrightarrow \text{PbCO}_3 + 2\text{NaNO}_3$ $\text{Pb(NO}_3)_2 + \text{K}_2\text{CO}_3 \longrightarrow \text{PbCO}_3 + 2\text{KNO}_3$
 - $(l) Pb(NO₃)₂ + Na₂SO₄ \longrightarrow PbSO₄ + 2NaNO₃$ $Pb(NO₃)₂ + H₂SO₄ \longrightarrow PbSO₄ + 2HNO₃$
 - (m) CuO + H₂SO₄(dil.) \longrightarrow CuSO₄ + H₂O CuCO₃+H₂SO₄(dil.) \longrightarrow CuSO₄+H₂O+ CO₂↑
 - (n) Mg + 2HCl(dil.) \longrightarrow MgCl₂ + H₂ \uparrow MgO + 2HCl(dil.) \longrightarrow MgCl₂ + H₂O MgCO₃+2HCl(dil.) \longrightarrow MgCl₂ + H₂O + CO₂ \uparrow
 - (o) $AgNO_3 + HCl(dil.) \longrightarrow AgCl + HNO_3$
 - Q19. Choosing only substances from the list given below in the box, write equations for the preparation of:
 - (i) Iron (II) sulphate

- (ii) Iron (II) chloride
- (iii) Iron (III) chloride
- (iv) Iron (II) sulphide
- (v) Zinc sulphate
- (vi) Copper sulphate
- (vii) Sodium chloride
- (viii) Copper carbonate

Iron, Copper,	dilute Sulphuric acid	Copper oxide
Sulphur, Chlorine,	dilute Hydrochloric acid	Sodium carbonate
Sodium, Zinc	dilute Nitric acid	Sodium hydroxide

- Ans. (i) Fe + $H_2SO_4 \longrightarrow FeSO_4 + H_2 \uparrow$
 - (ii) Fe + 2HCl \longrightarrow FeCl₂ + H₂ \uparrow
 - (iii) $2\text{Fe} + 3\text{Cl}_2 \longrightarrow 2\text{FeCl}_3$
 - (iv) Fe + S $\xrightarrow{\Delta}$ FeS
 - (v) Zn + H₂SO₄ \longrightarrow ZnSO₄ + H₂ \uparrow
 - (vi) CuO + H₂SO₄ \longrightarrow CuSO₄ + H₂O
 - (vii) NaOH + HCl --- NaCl + H2O
 - (viii) $CuO + H_2SO_4 \longrightarrow CuSO_4 + H_2O$ $CuSO_4 + Na_2CO_3 \longrightarrow CuCO_3 + Na_2SO_4$
- Q20. Starting from Lead nitrate how will you prepare the following named salts in laboratory. Write only the balanced chemical equations in support of your answer.
 - (i) Lead chloride
 - (ii) Lead sulphate
 - (iii) Lead carbonate
 - (iv) Lead sulphide
- Ans. (i) $Pb(NO_3)_2 + 2HCl(dil.) \longrightarrow PbCl_2 + 2HNO_3$
 - (ii) Pb(NO₃)₂ + Na₂SO₄ \longrightarrow PbSO₄ + 2NaNO₃
 - (iii) Pb(NO₃)₂ + Na₂CO₃ \longrightarrow PbCO₃ + 2NaNO₃
 - (iv) Pb(NO₃)₂ + Na₂S \longrightarrow PbS + 2NaNO₃
- Q21. What is the expected pH of the following solution
 - (i) which liberates carbon dioxide from metallic carbonates?
 - (ii) which liberates ammonia on heating with ammonium salts?
 - (iii) of pure water?

- Ans. (i) Less than seven
 - (ii) More than seven
 - (iii) Equal to seven.
- Q22. Solutions A, B, C and D are having pH 1, 5, 8 and 14 respectively. In this context answer the following questions.
 - (i) Choose the solutions which are acidic in nature.
 - (ii) Choose the solutions which are basic in nature.
 - (iii) Choose the solutions which are highly acidic and highly alkaline.
 - (iv) Choose the solutions which contain both ions and molecules.
- Ans. (i) A and B
 - (ii) C and D
 - (iii) Highly acidic A Highly alkaline - D
 - (iv) B and C
- Q23. Give one example each of the sodium salts as
 - (i) acid salt
 - (ii) normal salt
 - (iii) mixed salt
 - (iv) complex salt.
- **Ans.** (i) Sodium bisulphate or Sodium bicarbonate.
 - (ii) Sodium sulphate or Sodium carbonate.
 - (iii) Sodium potassium sulphate.
 - (iv) Sodium argentocyanide.
- Q24. What is the pH of the following solutions?
 - (i) Ferric chloride
 - (ii) Sodium acetate
 - (iii) Sodium chloride
- Ans. (i) Less than 7
 - (ii) More than 7
 - (iii) Equal to 7

- Q25. On adding pure water to acidic oxide and soluble basic oxide separately. How does the pH of pure water change?
- Ans. On adding pure water to acidic oxide the pH decreases. On adding pure water to soluble basic oxide the pH increases.
- Q26. Give only equations for the conversion of insoluble lead carbonate to lead chloride.
- Ans. $PbCO_3 + 2HNO_3 \longrightarrow Pb(NO_3)_2 + H_2O + CO_2$ $Pb(NO_3)_2 + 2HCl \longrightarrow PbCl_2 + 2HNO_3$

 $Pb(NO_3)_2 + 2NaCl \longrightarrow PbCl_2 + 2NaNO_3$

- Q27. Write balanced chemical equation to support each of the statement given below (use only dilute sulphuric acid).
 - (i) Basic oxide + Acid ------ Salt + Water
 - (ii) Amphoteric oxide + Acid ------ Salt +

O.H + COVO OF Water

- (iii) Alkali + Acid ------ Salt + Water
- (iv) Base + Acid ----- Salt + Water
- (v) Active metal + Acid \longrightarrow Salt +

Hydrogen

- (vi) Metallic carbonate + Acid ------ salt + Water + Carbon dioxide
- Ans. (i) MgO + $H_2SO_4 \longrightarrow MgSO_4 + H_2O$
 - (ii) $ZnO + H_2SO_4 \longrightarrow ZnSO_4 + H_2O$
 - (iii) $2NaOH + H_2SO_4 \longrightarrow Na_2SO_4 + 2H_2O$
 - (iv) $Cu(OH)_2 + H_2SO_4 \longrightarrow CuSO_4 + 2H_2O$
 - (v) $Mg + H_2SO_4 \longrightarrow MgSO_4 + H_2$
 - (vi) $Na_2CO_3 + H_2SO_4 \longrightarrow Na_2SO_4 + H_2O + CO_2$
- Q28. Name two salts each which an dissolving in water produces
 - (i) Acidic
- (ii) Basic
- (iii) Neutral solution
- (i) Copper sulphate and Ammonium chloride
 - (ii) Potassium acetate and Sodium sulphide
 - (iii) Sodium chloride and Potassium sulphate

the preparation of:

Let's Recall

ill You	ur Answ	er in t	he Space	Given for E	ach Quest	ion.						
Q1. M	latch the	e follo	wing:									•
A			olumn -I	ebine beloul			Column	ı-II	SW SISS			
4 3		(i) A	eid			(a)	Washing s	oda				
		(ii) B				(b)	Ferric chlo	oride				
			fflorescent	hate on breat			Hydroxyl i		and the same			
		A STATE OF THE PARTY OF THE PAR	eliquescer				Concentra		huric acı	d		
			ygroscopi			(e)	Hydrogen	ion	Mark - 1			
Ans.	(i)		(ii) ((iii)		(iv) ((v)			
	3.		Column -	ī			Colum	n-II				
	(middle) fro		erric chlo			(a)	Precipitat	ion				
			odium chl				Simple dis			THE R. C.		
				sulphate			Metal oxid		ilute acid	Pact,		- 1
The same			ead chlor			2.7%	Synthesis					
			opper sul			(e)	Neutraliz	ation				120
Ans.	(i) ((ii)		(iii)		(iv)		(v)	O ave 1		3.6
	c.		Column .	I			Colum	n-II				
Aus			otash alu			(a)	Acid salt					
			odium su			(b)) Double sa	alt				
				bisulphate		30.00) Basic sal					
		(iv) (Copper hy	droxychloride		(d) Normal s	alt			11112	
Ans.	(i)		(ii)		(iii)	175	(iv)					
	.,0				W With a							
Q2.	Fill in the	he bla	nks.	l whorens gol	huma in ov	vaen t	o form		w	hich on	dissol	ving
le for	(i) Ele	ement '	A' is a non	metal and it	which	turn	s blue litm					
		water p	roduces_	metals react	with dilute	acids	to liberate				160	
	(ii)				WIGH GHADE	sm		St				
			lioxide is l		mds (6)		se.					
	(iv) An	nmoniu	m hydrox	ide is a	estil (b)		whereas in	soluble s	alts are	generally	prep	ared
			alts are pr	epared by			Wilercas in		177	G	100	
Am	by			·	called as			(0)				
	(vi) Hy	ygrosco	pic substa	nces are also	loured inco	luble	salt.	Stand at				
	(vii)			is a green co	of motton of	ac	attomine.					
	(viii)	1 10 9110	7 707	has a smell	of rotten eg	go.	turn lime	water m	ilkv.			
	(ix) Bo	oth		and			_ turn inne	100				
	(x) Or	n addin	g acid to	methyl orange	it changes	to	II. Muliperi	ig la Bol			(131)	
Ω3.	State v	whethe	r the foll	owing state	ments are	True	or False.		1 93			
A 16				neutral oxide.				17				
4 1												
			is a triba						leidnes.			
	(iii) pH	of pure	water is	The state of the same	lata (n).		Drivigatio					
	(iv) Anl	hydrou	s sodium o	carbonate is ef	florescent.							
	(v) Ble	aching	powder is	a mixed salt.								
	1000		The same of the sa									

(1	(i) Sodium chioride is	double sait.				OTHER DESIGNATION OF THE PERSON OF THE PERSO
(v	ii) Ferro cyanide ion is	s a complex ion.				
(v	iii) Soluble bases are	called Alkalies.		ace Given for Each		dr mi restrictue de serrice che
(i.	x) Phosphoric acid is a	a weak dibasic acid.				Watch'thefollow
(.	x) Non-metals react b	oth with dilute as well	as cor	centrated acids.		other aridic medicable
Q4. (Complete and balance	ce the following cher	nical	equations (all reac	ctant	s are given).
						Single (in) Bush
		Room temperature				
		△ → +		100,000		
	(v) Fe + Cl ₂ \longrightarrow _	Illemitel +_				
	$vi)$ S + HNO ₃ \longrightarrow			To Control		
	Annual An	rea shizo lesseM (e)	10	17 We beschildingsmir	abog	Michell Propertion to
						stal (with mi given helf-y
		→(vi)+1				political materia.
		N-mmulo0 +				
		+ (a) Acid salt				
		→ +			mon	AND THE PERSON NAMED IN
		——————————————————————————————————————				
			Plant of Control Control Control			
		→+				
	Call State Manufacture 17		ich or	alv one option is co	rrec	t. Dark the bubble for
c	orrect answer.	oteracid of a block	312399	- Jone option is co		Dark the bubble lor
(i) Phosphorous acid is					
	(a) weak tribasic aci	- N N N N N N N N N N N N N N N N N N N	(b)	strong dibasic acid		amelian make Auto
	(c) weak dibasic acid	dulgeni enorodw	(d)	strong tribasic acid		
Ans.	(a)	b	C	onder March 1973		
(i	i) Ammonium salts on	heating with alkali pr	oduce	S		TO THE OWNER OF THE
	(a) hydrogen	(b) ammonia	(c)	nitrogen	(d)	None of these.
Ans.		(b) small man	(c)	Jo Design Brooks	(d)	died (set) Both
(ii	i) The aqueous solutio	n of potassium chloride	is		0	
	(a) alkaline	(b) neutral	(c)	acidic	(d)	amphoteric.
Ans.	(a)	(b)	(0)		(d)	Shade In Carrie (4)
(it) All acids essentially				0	
8	(a) oxygen	(b) nitrogen	(c)	carbon	(d)	hydrogen.
Ans.	(a)	(b)	(c)	STATUTE OF STATEOUS STATE	3312523	oq gaidasəldi (u)
70 0	- 1 ® CI				0	
70 7	ogether with ® Chemis	try (ICSE)-X				

(v)	Choose hydro-acid fro	om the following acids.		S AND THE REST OF		
((a) Acetic acid	(b) Carbonic acid	(c)	Hydrochloric acid	(d)	Sulphuric acid.
ns.	(a)	(b)	(c)		d	
(vi)	Choose oxy-acid from	the following acids.	Syringe in			0.69.5
	(a) Hydrochloric acid	(b) Hydroiodic acid	(c)	Nitric acid	(d)	Hydrofluoric acid.
ns.	(a)	(b)	0		(d)	
(vii)	Hydrated copper sul	phate on heating chang	ges to			(0) neutralization
	(a) blue and crystalli	ine	(b)	white and crystalli		
	(c) blue and amorphe	ous the shit shixab no	(d)	white and amorph	ous.	ign Pant on Pinte
ns.	(a)	b	0		(d)	
(vii	i) Iron (II) sulphide is	s prepared in laborator	y by		Hul.	
	(a) neutralization	(b) precipitation	(c)	synthesis	(d)	None of these.
lns.	(a)	b	C		(d)	N. Bes. R. L. Towns Co.
(ix)	The acid anhydride	of sulphuric acid is		DURA HM + ID-M		
	(a) sulphur dioxide	AND THE RESERVE	(b)	sulphur trioxide		
	(c) sulphur monochl	loride	(d)	None of these.	0	F (u) 2Fe+3CL,
Ans.	(a)	b	C	LONG FOREST OF	d	(0)(H9 + S ()a)
(x)		d softum aniphate (Sa	(2)	colourless and odd	nurlas	e gas
	(a) greenish yellow		(b) (d)	None of these.	Juries	s gas
	(c) reddish brown co	oloured gas		of tentesting to a	a	Least and the second
Ans.	(a)	1 -barres solution 'B'	bac n	H-4 In this context	choos	e the correct statement
(xi		d in comparison to B	(b)		e wea	ak bases
	(c) 'A' is a strong ac			'A' is strong base		
		(b)	(c	AICL + SH,O Inc)	(d	Lain Al O + EHCI -
Ans.	ii) The pH of copper cl		halas	Fr. William Belly	the th	Print Polices pull thirty on
(36)	(a) mole than 7	(b) Less than 7	(c)	Equal to 7	(d)	None of these
		(h)	(c	The HCL eller	(d	20, Na,CO, NaCE
Ans.	iii) In laboratory con	per sulphate is prepare	ed by	mous sulphate	_	
(x	(a) displacement	(b) Synthesis	(c)	Direct combination	on (d) Decomposition by acid
	and the second second		Co	r chloride	(d	
Ans.	(a)	rectly converts phospho		ol eschanata	_	
(30	(a) Dilute sulphuri		(b)		l.	
	(c) Dilute nitric ac		(d) Concentrated nit	ric ac	eid
			(0		(d	
Ans.	Soluble salt like so	odium sulphate can be p			the p	rocess of
(2	(a) Titration	(b) Synthesis	(c) Displacement	(d	Decomposition
			((d	
Ans.	(a)	6	0		_	To the later of

ANSWERS (iii) a (iv) b (ii) c **1. A.** (*i*) *e* (v) d(iii) b (iv) a $\mathbf{B}.$ (i) d(ii) e (iii) a (ii) d C. (i) b (iv) c 2. (i) acidic oxides, acids, red (ii) reactive, hydrogen (iii) burning sulphur (iv) weak (v) neutralization reactions, precipitation reactions (vi) drying agents (vii) copper carbonate (viii) Hydrogen sulphide (ix) Carbon dioxide and Sulphur dioxide (x) Red or Pink **3.** (*i*) False (ii) False (iii) True (iv) False (v) True (vi) False (vii) True (viii) True (ix) True (x) False 4. (i) $C + 4HNO_3 \longrightarrow CO_2 + 2H_2O + 4NO_2$ (ii) NaCl + H2SO4 Room temperature NaHSO4 + HCl (iii) $NH_4Cl + NaOH \xrightarrow{\Delta} NaCl + NH_3 + H_2O$ ebiznib nudglun (a) (iv) 2Pb(NO₃)₂ $\xrightarrow{\Delta}$ 2PbO + 4NO₂ + O₂ (v) $2\text{Fe} + 3\text{Cl}_2 \longrightarrow 2\text{FeCl}_3$ (vi) S + 6HNO₃ \longrightarrow H₂SO₄ + 2H₂O + 6NO₂ (vii) FeCl₃ + 3NH₄OH \longrightarrow Fe(OH)₃ + 3NH₄Cl (viii) 3Fe + 4H₂O \Longrightarrow Fe₃O₄ + 4H₉ (ix) Na₂S + H₂SO₄ \longrightarrow Na₂SO₄ + H₂S (x) $Na_2SO_3 + 2HCl \longrightarrow 2NaCl + H_2O + SO_2$ (xi) Na₂CO₃ + H₂SO₄ \longrightarrow Na₂SO₄ + H₂O + CO₅ (xii) Pb(NO₃)₂ + 2HCl \longrightarrow PbCl₂ + 2HNO₃

(xiv) Al₂O₃ + 6HCl \longrightarrow 2AlCl₃ + 3H₂O (xv) 2KOH + H₂SO₄ \longrightarrow K₂SO₄ + 2H₂O

(viii) c (ix) b (x) c (xi) c (xii) b (xiii) d (xiv) d (xv) a

(iv) d (v) c (vi) c (vii) d

(xiii) 2Al + 2NaOH + 2H₂O \longrightarrow 2NaAlO₂ + 3H₂

(ii) b (iii) b

(x) Zinc sulphide

(ix) Copper carbonate