
Astatilial Chemistry

REFERENCE BOOK



LATEST SYLLABUS

NOW IN NEW - FOUR COLOUR ON GLOSS PAPER AND IN HARD-BOUND JOURNAL FORMAT

# LATEST SYLLABUS FOR STD, IX - PRACTICAL CHEMISTRY

#### INTERNAL ASSESSMENT OF PRACTICAL WORK

Candidates will be asked to observe to the effect of reagents and/or of heat on substances supplied to them. The exercises will be simple and may include the recognition and identification of certain gases listed below.

Gases: Hydrogen, Oxygen, Carbon dioxide, Chlorine, Hydrogen chloride, Sulphur dioxide, Hydrogen sulphide, Ammonia, Water vapour, Nitrogen dioxide.

Candidates are expected to have completed the following minimum practical work.

#### SIMPLE EXPERIMENTS ON:

- 1. Heat the given (unknown) substance, make observations Identify any products and make deductions where possible.
  - (a) copper carbonate, zinc carbonate (b
- (b) washing soda, copper sulphate crystals
  - (c) zinc nitrate, copper nitrate, lead nitrate (d) ammonium chloride, iodine, ammonium dichromate
- 2. Add dilute sulphuric acid to the unknown substance -
  - Warm if necessary, make observation, identify the product and make deductions.
- (a) a sulphide
- (b) a carbonate
- (c) a metal
- 3. Apply the flame test to identify the metal in the unknown substance.
  - (a) a sodium salt
- (b) a potassium salt
- (c) a calcium compound
- 4. The percentage composition of a mixture of powdered salt and water-washed sand.

  The experiment would test techniques in dissolving, filtering or decanting, washing and weighing. It may be counted out as taking too much time. The weakness could be met by supplying a given weight of the mixture; also by choosing sand of such grain size that filtering or decanting will not be slow and yet not so large that separation of salt and sand cannot be done simply by sorting out mechanically the sand from the salt. The experiment should take about 20 minutes using
- 10g mixture (4g sand, 6g salt).
   5. Simple experiments based on hard water and soft water identification of hardness simple softening by heating the temporary hard water,
  - using washing soda and advantage of using detergents over soap in hard water.
- 6. Find out the sources of pollution of water bodies in the locality and determine the quality of water.

# PRACTICAL CHEMISTRY CHART

I. IDENTIFICATION	ON OF GASES
NEUTRAL GASES	
1. WATER VAPOUR	LITMUS TEST: Neutral to litmus.
	TESTS FOR THE GAS:
Colour: Colourless	• Turns white anhydrous copper sulphate - blue.
Odour: Odourless	$CuSO_4$ [white] + $5H_2O \rightarrow CuSO_4 \cdot 5H_2O$ [blue]
	• Turns blue cobalt chloride paper – pink.
	$CoCl_2$ [blue] + $2H_2O \rightarrow CoCl_2 \cdot 2H_2O$ [pink]
2. HYDROGEN [H <sub>2</sub> ]	LITMUS TEST: Neutral to litmus.
	TESTS FOR THE GAS:
Colour: Colourless	Burning wooden splinter is- extinguished in hydrogen.
Odour: Odourless	• Hydrogen burns with a – pale blue flame producing a 'pop' sound.
	$2H_2 + O_2 \rightarrow 2H_2O$ [on kindling]
3. OXYGEN [O <sub>2</sub> ]	LITMUS TEST: Neutral to litmus.
Colour: Colourless	* **TESTS FOR THE GAS:  • Rekindles – a glowing wooden splinter.
Odour: Odourless	Absorbed in colourless alkaline pyrogallol solution which turns - brown.
ACIDIC GASES	
4. CARBON DIOXIDE [CO <sub>2</sub> ]	LITMUS TEST: Moist blue litmus turns faint red.
	TESTS FOR THE GAS:
Colour: Colourless	Burning wooden splinter is – extinguished in carbon dioxide.
Odour: Odourless	On passage through lime water, it turns lime water – milky.
	$Ca(OH)_2 + CO_2 \rightarrow CaCO_3 \downarrow$ [white ppt insoluble] + H <sub>2</sub> O
	The <i>milkiness disappears</i> – on passage of excess carbon dioxide.
To office and administration of the second	$CaCO_3 + H_2O + CO_2 \rightarrow Ca(HCO_3)_2$ [soluble]  The gas has no effect on acidified KMpO or K Cr O solution
5 SUI PHUR DIOVIDE ISO 1	• The gas has <i>no effect</i> on acidified KMnO <sub>4</sub> or K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub> solution. <b>LITMUS TEST:</b> Moist blue litmus turns red.
J. SCETTICK DIOXIDE [SO <sub>2</sub> ]	TESTS FOR THE GAS:
Colour: Colourless	Burning wooden splinter is – <i>extinguished</i> in sulphur dioxide.
Odour: Suffocating	On passage through lime water, it turns lime water - milky.
	$Ca(OH)_2 + SO_2 \rightarrow CaSO_3 \downarrow $ [white ppt insoluble] + H <sub>2</sub> O
	The milkiness disappears - on passage of excess sulphur dioxide.
	$CaSO_3 + H_2O + SO_2 \rightarrow Ca(HSO_3)_2$ [soluble]
STATE OF THE STATE	• Turns acidified potassium permanganate from - pink to clear colourless.
	$2KMnO_4 + 2H_2O + 5SO_2 \rightarrow K_2SO_4 + 2MnSO_4 + 2H_2SO_4$
	• Turns acidified potassium dichromate from - orange to clear green.
	$K_2Cr_2O_7 + H_2SO_4 + 3SO_2 \rightarrow K_2SO_4 + Cr_2(SO_4)_3 + H_2O$

ACIDIC GASES [Contd.]	The season of the constitution of the season
6. CHLORINE [Cl <sub>2</sub> ]	LITMUS TEST: Moist blue litmus turns red and then gets bleached.
The state of the s	TESTS FOR THE GAS:
Colour: Greenish yellow	• Turns moist blue litmus red and then - bleaches it.
Odour: Pungent	$Cl_2 + H_2O \rightarrow HCl + HOCl; HOCl \rightarrow HCl + [O] [nascent]$
LOBOL MINE	Colouring matter [litmus] + [O] → Colourless or bleached product
HO, LIVE SAVE OF IN	Turns moist starch iodide paper [KI + starch solution] - blue black.
Tylidd bank bledt a.e.	$Cl_2 + 2KI \rightarrow 2KCl + I_2$
	Starch + I <sub>2</sub> → Blue black colour
7. HYDROGEN CHLORIDE [HCl]	LITMUS TEST: Moist blue litmus turns red.
	TESTS FOR THE GAS:
Colour: Colourless Odour: Pungent	Gives dense white fumes when a glass rod dipped in ammonia solution is brought near hydrochloric acid vapours.
Out our Fungerio	NH <sub>3</sub> + HCl [vapours] → NH <sub>4</sub> Cl [dense white fumes]
	• Forms a curdy white precipitate – on passage through AgNO <sub>3</sub> solution.
1 2 2 2 2 4 1 2	AgNO <sub>3</sub> [aq.] + HCl → AgCl ↓ [curdy white ppt.] + HNO <sub>3</sub>
NO DEDUCTION -	The precipitate of AgCl is <i>soluble</i> in NH <sub>4</sub> OH but <i>insoluble</i> in dil. HNO <sub>3</sub> .
8. HYDROGEN SULPHIDE [H <sub>2</sub> S]	LITMUS TEST: Moist blue litmus turns red.
I SHARE THE WAY	TESTS FOR THE GAS:
Colour: Colourless	• Turns acidified potassium permanganate from - pink to colourless.
Odour: Rotten egg	$2KMnO_4 + 3H_2SO_4 + 5H_2S \rightarrow K_2SO_4 + 2MnSO_4 + 8H_2O + 5S$
william warning restrict	• Turns acidified potassium dichromate from - orange to green.
	$K_2Cr_2O_7 + 4H_2SO_4 + 3H_2S \rightarrow K_2SO_4 + Cr_2(SO_4)_3 + 7H_2O + 3S$
consider contentals	The above tests are answered also by sulphur dioxide [SO <sub>2</sub> ], but in case of H <sub>2</sub> S, <i>yellow particles of sulphur</i> are seen and the solution – is hence not clear colourless or clear green respectively.
	Turns moist lead acetate paper - silvery black.
The state of the state of	Pb(CH <sub>3</sub> COO) <sub>2</sub> [colourless] + H <sub>2</sub> S → PbS↓ [black] + 2CH <sub>3</sub> COOH
9. NITROGEN DIOXIDE [NO	LITMUS TEST: Moist blue litmus turns red.
Series of South Series	TESTS FOR THE GAS:
Colour: Reddish brow	• It liberates iodine [violet vapours] from potassium iodide [KI] solution.
Odour: Irritating	$2KI + 2NO_2 \rightarrow 2KNO_2 + I_2$ [Turns potassium iodide paper brown]
boundt.	• It dissolves in cold water to give a mixture of nitrous and nitric acid in solution which acts on - blue litmus and turns it red.
Salarichus suiz si redom	2NO <sub>2</sub> + H <sub>2</sub> O → HNO <sub>2</sub> + HNO <sub>3</sub> [hence NO <sub>2</sub> is called a mixed acid anhydride]

<ul> <li>10. AMMONIA [NH<sub>3</sub>]</li> <li>LITMUS TEST: Moist red litmus turns blue.</li> <li>TESTS FOR THE GAS:</li> <li>Gives dense white fumes when a glass rod – dipped in conc. HCl acid is brought near ammonia gas.</li> <li>NH<sub>3</sub> + HCl [conc.] → NH<sub>4</sub>Cl [dense white fumes]</li> <li>On passage through copper [II] sulphate solution [CuSO<sub>4</sub>] – it gives a pale blue precipitate. [NH<sub>3</sub> dissolved in H<sub>2</sub>O gives NH<sub>4</sub>O CuSO<sub>4</sub> + 2NH<sub>4</sub>OH → (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> + Cu(OH)<sub>2</sub> pale blue ppt.]</li> <li>On passage of excess ammonia [or excess NH<sub>4</sub>OH solution] to pale blue precipitate dissolves to give a deep blue or inky bits</li> </ul>								
<ul> <li>Colour: Colourless Odour: Pungent</li> <li>Gives dense white fumes when a glass rod – dipped in conc. HCl acid is brought near ammonia gas.  NH<sub>3</sub> + HCl [conc.] → NH<sub>4</sub>Cl [dense white fumes]</li> <li>On passage through copper [II] sulphate solution [CuSO<sub>4</sub>] – it gives a pale blue precipitate. [NH<sub>3</sub> dissolved in H<sub>2</sub>O gives NH<sub>4</sub>O  CuSO<sub>4</sub> + 2NH<sub>4</sub>OH → (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> + Cu(OH)<sub>2</sub> ↓ [pale blue ppt.] On passage of excess ammonia [or excess NH<sub>4</sub>OH solution] to pale blue precipitate dissolves to give a deen blue or inku bit</li> </ul>								
<ul> <li>Odour: Pungent</li> <li>dipped in conc. HCl acid is brought near ammonia gas.</li> <li>NH<sub>3</sub> + HCl [conc.] → NH<sub>4</sub>Cl [dense white fumes]</li> <li>On passage through copper [II] sulphate solution [CuSO<sub>4</sub>] – it gives a pale blue precipitate. [NH<sub>3</sub> dissolved in H<sub>2</sub>O gives NH<sub>4</sub>O</li> <li>CuSO<sub>4</sub> + 2NH<sub>4</sub>OH → (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> + Cu(OH)<sub>2</sub> √ [pale blue ppt.]</li> <li>On passage of excess ammonia [or excess NH<sub>4</sub>OH solution] to pale blue precipitate dissolves to give a deep blue or inku bit</li> </ul>								
<ul> <li>On passage through copper [II] sulphate solution [CuSO<sub>4</sub>] - it gives a pale blue precipitate. [NH<sub>3</sub> dissolved in H<sub>2</sub>O gives NH<sub>4</sub>O CuSO<sub>4</sub> + 2NH<sub>4</sub>OH → (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> + Cu(OH)<sub>2</sub>  [pale blue ppt.] On passage of excess ammonia [or excess NH<sub>4</sub>OH solution] to pale blue precipitate dissolves to give a deep blue or inku bit</li> </ul>								
it gives a pale blue precipitate. $[NH_3 \text{ dissolved in } H_2O \text{ gives } NH_4O  CuSO_4 + 2NH_4OH \rightarrow (NH_4)_2SO_4 + Cu(OH)_2 \downarrow [pale blue ppt.]$ On passage of excess ammonia [or excess $NH_4OH \text{ solution}]$ to pale blue precipitate dissolves to give a deep blue or inky by								
On passage of excess ammonia [or excess NH <sub>4</sub> OH solution] to pale blue precipitate dissolves to give a deep blue or inky by								
pale blue precipitate dissolves to give a deen blue or inky bi								
solution of the complex salt – tetramine copper [II] sulphate.								
$Cu(OH)_2 + (NH_4)_2SO_4 + 2NH_4OH \rightarrow [Cu(NH_3)_4]SO_4 + 4H_2$								
[pale blue [tetramine ppt.] copper (II) sulphate]								
Turns Nessler's reagent from - colourless to pale brown or gives reddish brown ppt. [iodide of Millon's base] on passage of NH <sub>3</sub> in exce								
Nessler's reagent is an alkaline soln. of potassium mercuric iodide [K <sub>2</sub> Hg								
II ACTION OF HEAT - On the given [unknown] substance								
SUBSTANCE PRODUCTS OBSERVATION AND DEDUCTION								
1. Copper carbonate • Original colour - Light green								
Colour change - Turns black on strong heating								
CuCO <sub>3</sub> → CuO + CO <sub>2</sub> [g] • Gas evolved - Carbon dioxide								
[Copper carbonate] [Copper oxide] [Carbon a] Colour & odour - Colourless, odourless [light green] [black] dioxide] b] Nature - Slightly acidic to litmus								
c] Test - Lime water turns mill [KMnO <sub>4</sub> -no effect].								
• Residue - Black copper oxide formed.  Deductions:  The light green powder is copper carbona.								
2. Zinc carbonate • Original colour - White								
ZnCO <sub>3</sub> → ZnO + CO <sub>2</sub> [g] • Colour change - Turns yellow on heatin								
Live 3								
[Zinc carbonate] [Zinc oxide] [Carbon • Gas evolved - Carbon dioxide								

c] Test - Lime water turns milky

Deductions: The white powder is zinc carbonate.

• Residue - Zinc oxide formed.

[KMnO<sub>4</sub>-no effect].

[yellow-hot, white-cold]

SUBS	TANCE	PRODUCTS	OBSERVATION AND DEDUCTION
[Wash	ng soda 3.10H2O → ing soda] white]	Na <sub>2</sub> CO <sub>3</sub> .H <sub>2</sub> O+ 9H <sub>2</sub> O [Sodium [water carbonate] vapour] [white]	<ul> <li>Original colour - White</li> <li>Colour change - Remains white on heating.</li> <li>Gas evolved - Water vapour [water of crystallization] a] Colour - Colourless liquid b] Nature - Neutral to litmus c] Test - Turns cobalt chloride blue to pink.</li> <li>Residue - White sodium carbonate is formed.</li> <li>Deductions: The white powder is washing soda.</li> </ul>
CuSO	r sulphate  1.5H <sub>2</sub> O → r sulphate] s - blue]	CuSO <sub>4</sub> + 5H <sub>2</sub> O [Copper [water sulphate] vapour] [anhydrous-white]	<ul> <li>Original colour - Blue [hydrous]</li> <li>Colour change - Turns white [anhydrous] on heating. On strong heating a black residue is formed.</li> <li>Gas evolved - [Water vapour [initial]] - SO<sub>2</sub>&amp;O<sub>2</sub> Sulphur dioxide - colourless, acidic to litmus, turns KMnO<sub>4</sub> soln. pink to colourless.</li> </ul>
20	ong heating CuSO <sub>4</sub> →	2CuO + 2SO <sub>2</sub> + O <sub>2</sub> [Copper [Sulphur [Oxygen oxide] dioxide] [black]	Oxygen - colourless, odourless, neutral to litmus, relits glowing splint.  • Residue-Initial-anhydrous copper sulphate, On strong heating - copper oxide.  Deductions: The blue powder is copper sulphate.  • Original colour - White
2Zr	nitrate  rong heating n(NO <sub>3</sub> ) <sub>2</sub>	2ZnO + 4NO <sub>2</sub> +O <sub>2</sub> [yellow-hot] [nitrogen [white-cold] dioxide]	<ul> <li>Colour change - Turns yellow on strong heating.</li> <li>Gasevolved - [Water vapour [initial]] NO<sub>2</sub> &amp; O<sub>2</sub> Nitrogen dioxide - reddish brown, acidic to litmus, turns acidified ferrous sulphate solution brown.</li> <li>Oxygen - colourless, neutral, relits glowing splint.</li> <li>Residue - Zinc oxide - formed [Yellow-hot, white-cold]</li> <li>Deductions: The white powder is hydrated zinc nitrate.</li> </ul>
2C	er nitrate (u(NO <sub>3</sub> ) <sub>2</sub> per nitrate) [blue]	2CuO + 4NO <sub>2</sub> + O <sub>2</sub> [Copper [Nitrogen [Oxygen oxide] dioxide] [black]	<ul> <li>Original colour - Blue</li> <li>Colour change - Turns black on heating.</li> <li>Gas evolved - [Water vapour [initial]] NO<sub>2</sub>&amp;O<sub>2</sub> <i>Nitrogen dioxide</i> - reddish brown, acidic to litmus, turns acidified ferrous sulphate soln. brown.         <i>Oxygen</i> - colourless, neutral, relits glowing splint.         <ul> <li>Residue - Black copper oxide is formed.</li> <li>Deductions: The blue powder is copper nitrate.</li> </ul> </li> </ul>

	SUBSTANCE	PRODUCTS	OBSERVATION AND DEDUCTION					
7.	Lead nitrate		Original colour - White crystalline solid					
	[lead nitrate]	2PbO + 4NO <sub>2</sub> + O <sub>2</sub> [litharge] [nitrogen [Oxygen]	• Colour change - Turns yellow on heating, decrepitates and melts.					
	[white]	[yellow] dioxide]	• Gas evolved - [Water vapour [initial]] NO <sub>2</sub> &O <sub>2</sub> Nitrogen dioxide - reddish brown, acidic to litmus, turns acidified ferrous sulphate solution brown.					
			Oxygen - colourless, neutral, relits glowing splint.					
			• Residue - Litharge is formed on strong heating which fuses with the glass.					
		THE PROPERTY OF THE PARTY OF TH	Deductions: The white powder is lead nitrate.					
8.	Ammonium chloride	antibot Chuldrent   home	Original colour - White crystalline					
	NH <sub>4</sub> Cl heat cool [Ammonium chloride] [white]	NH <sub>3</sub> + HCl  [Ammonia] [Hydrogen chloride] [colourless]	chloride sublimate which condenses on the					
		tribuni guran artir i	Residue - No residue is left in the test tube.					
		resolve Innight of	Deductions: The white powder is ammonium chloride.					
9.	Iodine	TRUMP TOUR PARTY.	Original colour - Violet crystals.					
	$I_2[s] \rightarrow$ [Iodine - solid] [violet]	2 I [vapours] [Iodine - vapours] [violet vapours]	<ul> <li>Colour change - Sublimes on heating evolving violet vapours.</li> <li>Gas evolved - Iodine vapours</li> </ul>					
		THE PARTY OF THE P	a] Colour - Violet vapours					
		relifes answers	b] Test-Fumes turns silver nitrate paper yellow					
	websyl bussil able	and we be the second	Residue - No residue is left in the test tube.					
10	A		Deductions: The violet crystals are of iodine.					
10.	Ammonium dichromate	and extending the land	Original colour - Orange					
	(NH <sub>4</sub> ) <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub> → [Ammonium dichromate] [orange]	Cr <sub>2</sub> O <sub>3</sub> + 4H <sub>2</sub> O + N <sub>2</sub> [Chromic oxide] [green]	On strong heating - decomposes violently with flashes of light leaving a voluminous green residue.					
	delica anomal batter	Sharest majornal	• Gases evolved - [Water vapour [initial]], N <sub>2</sub>					
	times within lieutenase	Selm brown	• Residue - Green chromic oxide on strong heating.					
	copper axide is thrive e-polyder is apper with	F Residue - Black Deductions: The bi	Deductions: The orange powder is ammonium dichromate.					

III ADDITION OF DIL. SULPHURIC ACID - 10 the given funktiowill substance									
SUBSTANCE									
1. SULPHIDE	<ul> <li>Colourless gas is evolved with a rotten egg smell.</li> <li>Gas turns lead acetate paper silvery black</li> </ul>	Gas : Hydrogen sulphide Substance: A sulphide Equation : Na <sub>2</sub> S + H <sub>2</sub> SO <sub>4</sub> → Na <sub>2</sub> SO <sub>4</sub> + H <sub>2</sub> S							
2. CARBONATE	<ul> <li>Colourless, odourless gas</li> <li>Gas turns lime water milky but has no effect on potassium permanganate solution.</li> </ul>	Gas : Carbon dioxide Substance: Carbonate or hydrogen carbonate Equation : $Na_2CO_3 + H_2SO_4 \rightarrow Na_2SO_4 + H_2O + CO_2$ $2NaHCO_3 + H_2SO_4 \rightarrow Na_2SO_4 + H_2O + 2CO_2$							
3. METAL	<ul> <li>Colourless, odourless gas is evolved with brisk effervescence</li> <li>Gas burns with a 'pop' sound when burning splint brought near it.</li> </ul>	Gas : Hydrogen Substance: Active metal Equation : $Mg + H_2SO_4 \rightarrow MgSO_4 + H_2$							
IV APPLIC	CATION OF FLAME TEST	- To identify	the metal in	the substance					
		Colour imparted to the flame	CO 200 100 100 100 100 100 100 100 100 100	Metallic radical					
and then h	num wire is thoroughly cleaned neated – uminous flame of a burner.	Golden yellow Pale yellow Sodium [Na <sup>1</sup>							
• The wire in non-lumin	wire imparts no colour it is conc. HCl and then- abstance to be identified. as then reintroduced into the - nous flame and the -	Lilac Violet Potassium [K <sup>1</sup> -  Brick red Pale green Calcium [Ca <sup>2+</sup>							
colour im	parted is noted.								
V DETER	MINATION OF % COMPC	SITION OF -	MIXTURE						
The state of the state of	METHOD	The Parks of the P	CALCULATIO	NS .					
powdere taken in	aixture of – ed salt & water wash sand is – a beaker and weighed.	Mass of beaker $= X g$ .  Mass of beaker $+ mixture = Y g$ .  Mass of mixture $= [Y - X] g$ $= [Y - X] g$							
to dissol Collect t separate		∴[Y – X] g	Mass of washed & dried sand = $Zg$ . $\therefore [Y - X] g \text{ of mixture contains } Zg \text{ of sand}$ $\therefore 100 g \text{ of mixture contains } \frac{Z \times 100}{[Y - X]} = \text{'A'}$						
• Weigh t	he washed and dried sand.	∴ A = Percentage of sand in mixture.							

# VI EXPERIMENTS BASED ON - Hard & Soft water

Hard water - is water which does not lather with ordinary soap & contains dissolved -Ca & Mg [bicarbonates, sulphates, chlorides]. Soft water - is water which lathers readily with ordinary soap & does not contain dissolved calcium and magnesium salts.

Hard water is further subdivided into - a] Temporary b] Permanent hard water

- Temporary hard water contains Ca and Mg bicarbonates in water.
- Permanent hard water contains Ca & Mg chlorides & sulphates in water.

# Experiment - Method

#### Observation - Result

#### **EXPERIMENT - I**

#### Differentiating hard water from soft water

- Two unknown samples 'X' and 'Y' containing hard water and soft water are taken separately in a trough or beaker.
- Ordinary soap is rubbed by the hands inside each sample.

#### EXPERIMENT - II

### Differentiating temporary & permanent hard water

- Two unknown samples 'A' and 'B' containing temporary & permanent hard water are taken- • The sample of water 'B' - does not lather with soap. separately in a trough or beaker.
- The water is boiled slowly, gases allowed to The boiled & filtered sample 'A' which lathers is escape out, and then the water is filtered.
- Ordinary soap is rubbed by the hands inside each filtered sample.

#### EXPERIMENT - III

# Temporary hard water softened by heating

- Temporary hard water is taken in a beaker and The boiled and filtered sample of heated slowly.
- · After the gases escape out, the water is filtered Result through a filter paper.
- Ordinary soap is rubbed inside the filtered solution.

#### **EXPERIMENT - IV**

#### Temporary and permanent hard water softened Observation by addition of washing soda

- Temporary & permanent hard water are taken separately in beakers & washing soda is added Result to each sample of water. The above solutions • Temporary hard water & Permanent hard water are filtered - to remove the precipitate formed.
- Ordinary soap is rubbed inside the filtered solution.

### **EXPERIMENT - V**

## Advantage of using detergents over soap

- A sample of hard water 'X' is taken and ordinary soap is rubbed inside the water.
- Another sample of hard water 'Y' is taken and o Detergents form lather even with hard water, while detergent is rubbed inside the water. [Detergents are sodium salts of sulphonic acid and] do not form scum.]

#### Observation

- One sample of water 'X' lathers with soap.
- The sample of water 'Y' does not lather with soap. Result
- Sample 'X' which lathers is soft water.
- Sample 'Y' which does not lather is hard water.

#### Observation

- One sample of water 'A' lathers with soap.
- Result
- temporary hard water whose hardness is removed by boiling. Sample 'B' is permanent hard water whose hardness cannot be removed by boiling.


#### Observation

- temporary hard water lathers readily with soap.
- Temporary hard water can be softened by heating.  $Ca(HCO_3)_2 \longrightarrow CaCO_3 \checkmark + CO_2 + H_2O$ [in temporary [ppt. filtered [filtered water hard water] out] is soft]


- The filtered sample of temporary & permanent hard water lathers - readily with ordinary soap.
- can be softened by using washing soda.


#### Observation

- · Lather forms in 'Y' but not in 'X'. Result
- ordinary soap is wasted due to formation of scum.  $Ca(HCO_3)_2 + 2NaSt \rightarrow Ca(St)_2 \downarrow + 2NaHCO_3$ [in hard water] [soap] SCHIM



# Voentitication of Gases






Dukillillialini of Ababa HYDROGEN CHLORIDE **GIVES DENSE WHITE FUMES WITH** GLASS ROD DIPPED IN AMMONIA SOLN. NaCI + Conc. H<sub>2</sub>SO<sub>4</sub> HYDROGEN SULPHIDE 8. **TURNS MOIST LEAD ACETATE PAPER** SILVERY BLACK METALLIC SULPHIDE + **DILUTE ACID** NITROGEN DIOXIDE 9. **REDDISH BROWN FUMES EVOLVED TURNS KI PAPER BROWN** COPPER + Conc. HNO<sub>3</sub> AMMONIA) 10. **TURNS MOIST GIVES DENSE WHITE RED LITMUS PAPER BLUE FUMES WITH GLASS ROD DIPPED IN** CONC. HCI NH4CI + **CALCIUM HYDROXIDE** 

# Action of heat on the given substance 3. LEAD [IV] OXIDE) 1. MERCURY [II] OXIDE 2. RED LEAD RELITS **RELITS** RELITS **GLOWING GLOWING GLOWING** SPLINT **SPLINT SPLINT** RESIDUE RESIDUE RESIDUE Yellow -Yellow -Silvery deposit fuses with glas fuses with g COPPER CARBONATE **MOIST BLUE LITMUS TURNS FAINT RED** RESIDUE ORIGINAL COLOUR **AFTER HEAT - Black Light Green**








# Action of heat on the given substance





# III. Flame Test



# VII DETERMINATION OF - The Quality of Water

Experiment - Find the sources of pollution of water bodies in the locality & determine the quality of water

- Sources of polluted water may include run off water from streets, garden lawns, septic tank
  discharges or waste water from sewage waste & other samples of fresh water & marine water.
- · The Water Quality Indicators include -

W	ater quality indicators	Test performed for - determining quality of water				
• Physical		Turbidity, temperature, specific conductance, dissolved solids - clarity, colour, odour, salinity etc.				
	Chemical	pH of water, dissolved oxygen in water, dissolved nutrients in water.				
	Biological	Presence of algae, bacteria, viruses etc.				

# Simple Tests - for determining the quality of a sample of water

Turbidity	Turbidity – refers to the particulate matter suspended in water bodies.  Measurement – By use of a handheld – turbidity meter which measures – the scattering of light by matter, when a beam of light is passed into water.  Conclusion – Measurement helps to calculate inputs from nutrients & soil erosion.
Temperature	Measurement - By use of an accurate thermometer.  Conclusion - Hot water from industrial plants, affects the - temperature of water bodies it enters.  Warm water holds less oxygen and hence affects survival of living species
Specific conductance	Measurement - Simple electrolytic cell - to determine the degree to which water can conduct electricity.  Conclusion - Low specific conductance indicates - less polluted water.
Dissolved solids	Measurement - The sample of water is filtered & then evaporated.  The solids left behind are then - chemically tested.  Conclusion - Dissolved inorganic matter such as - bicarbonate, chlorides macause hardness in water. Dissolved solids make water look & taste unpleasant.
pH of water	Measurement – By use of pH paper or electronic pH-meter.  Conclusion – Changes in pH of water affects dissolution of chemicals water and survival of certain organisms which cannot exist in highly acid water. [unpolluted water generally has a pH range around 6.5 to 8]
Dissolved oxygen	Measurement - The sample of water is tested by using special field kits  Conclusion - Unpolluted water have high amount of dissolved oxygen [Down whose presence & amount is responsible for survival of aquatic organism  [Temperature & speed of water affect the amount of dissolved oxygen]
Dissolved nutrients	Measurement - Residual water after evaporation is subjected to - chemical test Conclusion - Increase in nutrients affects - pH value, clarity & temperature of water. Increase in 'nitrogen & phosphorous' will result in eutrophication
 Bacteria	Measurement - Test for coliform bacteria in the water sample using various methods including a medium called m-endroth.  Conclusion - Presence of bacteria along with viruses are detrimental human health.

#### Additional Questions Give a chemical test to distinguish between the following gases: Hydrogen and oxygen Chlorine and nitrogen dioxide Carbon dioxide and sulphur dioxide Ammonia and hydrogen chloride e] Hydrogen chloride and hydrogen sulphide Sulphur dioxide and chlorine. f On heating which of the following substances i.e. copper carbonate, zinc carbonate, washing soda, copper sulphate, zinc nitrate, copper nitrate, lead nitrate, ammonium chloride and ammonium dichromate - relate to the reactions given below.

- - A white substance which leaves an amphoteric oxide as a residue [whose colour varies in the heated and in the cold state] and evolves a gas which turns lime water milky.
  - An efflorescent substance which leaves a residue having the same colour as the substance and evolves b] a gas which changes the colour of cobalt chloride paper.
  - A white solid which evolves two colourless gases which on cooling combine and condense on the cooler parts of the test tube.
  - A coloured substance which decomposes violently leaving a coloured residue and evolving two neutral gases one of which is unreactive or inert in nature.
  - A coloured substance which leaves a black residue and evolves two gases one of which is acidic and coloured and the other neutral and colourless.
  - A coloured substance which leaves on strong heating a black residue and evolves two colourless gases f] one of which is acidic and the other neutral.
  - A white crystalline solid which decrepitates on heating leaving a residue which fuses with the glass g and evolves two gases one of which is coloured and acidic.
  - An amorphous substance which turns from pale green to black on strong heating evolving a colourless, acidic gas as the only gaseous product.
- Give balanced equations for the following conversions affected by heat alone on the substances:
  - Copper carbonate to copper oxide; b] Hydrated copper sulphate to sulphur dioxide;
  - Ammonium dichromate to nitrogen; Copper nitrate to nitrogen dioxide; d]
  - Zinc carbonate to zinc oxide; f] Zinc nitrate to nitrogen dioxide
- Using dilute sulphuric acid how would you differentiate between:
  - b] Copper and magnesium. a] Sodium sulphide and sodium carbonate. How would you identify the gaseous products evolved.
- Using a platinum wire, conc. hydrochloric acid and a bunsen burner how would you distinguish between the three salts ie. sodium chloride, potassium chloride and calcium chloride. Explain in brief the method used for the same.
- A mixture consists of 20% sodium chloride and 80% sand. Explain practically in brief a simple method involved to ascertain the correct percentages in the mixture.
- Using given samples of temporary and permanent hard water, soft water, ordinary soap, detergent and washing soda how would you - a] distinguish between hard and soft water b] distinguish between temporary hard water and permanent hard water c] remove temporary hardness from water without using a chemical compound d] remove temporary hardness and permanent hardness from water using a chemical compound e] prove the advantage of detergent over soap.
- To determine the quality of water in different water samples collected from water sources, specific tests are performed. State four physical tests & three chemical tests performed to determine the quality of water. For each test explain how the test is conducted & the conclusion drawn about the quality of water.

0.1	Sele	ct the	correct gas	from A to	F which	match	es wi	th the descr	riptions	1 to 5	5.		[5]
2.1	A: (		B: SO,	C: NH <sub>3</sub>		ater va		E: Cl <sub>2</sub>	F: H				
	1.	- An	moist blue	9	and the	n bleac	hes it						
	2.		moist red										
	3.		lime water			mus pa	per sl	ightly pink					
	4.		cobalt chlo										
	5.		s lead aceta					lack.					
Q.2		ect the	correct sal						tion ex	hibits	the cha	nge in col	lour from [5]
		I - 1 to	0 5.				Lis	t-II					
	List		aroon to bl	ack				Copper car	bonate				
	1.	1000	green to bl		tal			Hydrated C			ate		
	2.		e to yellow	[neated sta	itej			Copper nit					
	3.		to black					Ammoniu		omate			
	4.		ge to green					Zinc nitrate					
	5.		to white										(E)
Q.3	Sta	te whi	ch of the s						s on the		decomp	osition.	[5]
	1.	Zinc	carbonate		shing soc	la 3.		ad nitrate	4,79	4.		nium dich	
	5.	Trile	ad tetroxide	e 6. Zino	nitrate	7	. M	ercury [II] o	oxide	8.	Annyar	ous coppe	er sulphate
0.4	4 Co	mplet	e the table	given belo	w.			nS <sub>O</sub> S	# Divis	OFF	A Portion	and the	[10]
	1.	- Ini	at on coppe		TEN T	• Col	our o	f acidic gas	evolve	d			
	2.		at on iodine			• Col	our o	f vapours e	volved				
	3.		at on ammo		romate	• Na	me of	neutral gas	evolve	ed	PART OF THE PART O	9,000,000	
	4.	He	at on coppe	r carbonate	e	• Tes	t for g	gas evolved					
	5.	22	at on zinc n			. Col	our o	f residue			RUERIK I		
	6.	. Ad	dition of di	1. H <sub>2</sub> SO <sub>4</sub> to	FeS	• Od	our o	f gas evolve	ed				
	7		dition of di			• Na	me of	residue ob	tained	W. W.	7 7 7		
	8		dition of di				t for	gas evolved	1	-	1000	TARK.	1175
	9	. He	at on sodiu	m chloride		• Co	lour i	mparted to	flame c	luring	flame to	est	
	1	0. Fla	me test for	calcium ch	loride	• Ac	id use	ed for flame	test				
0	E 60	lost th	e correct a	newer from	the wo	rds in l	rack	et.					[5]
Q			d and soft w						[dil	. acid	/ ordina	ry soap /	detergent].
	1.	Low	v specific co	nductance	of water	indica	tes	[less					
	<ul><li>2.</li><li>3.</li></ul>	The	type of w	ater softer	ed by a	ddition	of w	ashing soc	la is _	_ [p	ermanei	nt / tempo	orary / both
	1	type	oolluted wa	iter has		_ [high	/low	amount o	of dissol	ved o	xygen.		
	4.	The	sample of	hard wate	r which	lathers	with	soap after	boiling	& filt	ration co	ontains _	
	5.	[cal	cium chlori	de / calciu	m bicarb	onate].	Sate						

30 marks

Unit Test Paper - Practical Chemistry